MULTI-SIGHTLINE OBSERVATION OF NARROW ABSORPTION LINES IN LENSED QUASAR SDSS J1029+2623^{1,2}

Toru Misawa³, Cristian Saez^{4,5}, Jane C. Charlton⁶, Michael Eracleous^{6,7}, George Chartas⁸, Franz E. Bauer^{9,10,11}, Naohisa Inada¹², and Hisakazu Uchiyama¹³

A complete version

ABSTRACT

We exploit the widely-separated images of the lensed quasar SDSS J1029+2623 ($z_{\rm em}=2.197, \theta=$ 22".5) to observe its outflowing wind through two different sightlines. We present an analysis of three observations, including two with the Subaru telescope in 2010 February (Misawa et al. 2013) and 2014 April (Misawa et al. 2014b), separated by 4 years, and one with the Very Large Telescope, separated from the second Subaru observation by ~ 2 months. We detect 66 narrow absorption lines (NALs), of which 24 are classified as *intrinsic* NALs that are physically associated with the quasar based on partial coverage analysis. The velocities of intrinsic NALs appear to cluster around values of $v_{\rm ei} \sim 59,000, 43,000$, and $29,000 \ {\rm km \ s^{-1}}$, which is reminiscent of filamentary structures obtained by numerical simulations. There are no common intrinsic NALs at the same redshift along the two sightlines, implying that the transverse size of the NAL absorbers should be smaller than the sightline distance between two lensed images. In addition to the NALs with large ejection velocities of $v_{\rm ej} >$ 1,000 km s⁻¹, we also detect broader proximity absorption lines (PALs) at $z_{\rm abs} \sim z_{\rm em}$. The PALs are likely to arise in outflowing gas at a distance of $r \leq 620$ pc from the central black hole with an electron density of $n_{\rm e} \geq 8.7 \times 10^3$ cm⁻³. These limits are based on the assumption that the variability of the lines is due to recombination. We discuss the implications of these results on the three-dimensional structure of the outflow.

Subject headings: quasars: absorption lines – quasars: individual (SDSS J1029+2623)

1. INTRODUCTION

AGN outflows, potentially powered by one or more of a variety of mechanisms (e.g., radiation force, magnetic pressure, and magnetocentrifugal force), are important ingredients of quasar central engines and likely play a role in quasar and galaxy formation/evolution because: 1) they extract angular momentum from accretion disks allowing gas accretion to proceed (e.g., Blandford & Payne 1982; Emmering, Blandford, and Shlosman 1992; Konigl & Kartje 1994; Everett 2005), leading to the growth of

Electronic address: misawatr@shinshu-u.ac.jp

¹ Based on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

Based on observations obtained at the European Southern Observatory at La Silla, Chile in programs 092.B-0512(A)

- ³ School of General Education, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan
- ⁴ Korea Astronomy and Space Science Institute (KASI), 61-1, Hwaam-dong, Yuseong-gu, Deajeon 305-348, Republic of Korea
- ^b Department of Astronomy, University of Maryland, College

Park, MD 20742-2421 ⁶ Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802

⁷ Department of Astronomy & Astrophysics and Center for Gravitational Wave Physics, The Pennsylvania State University, 525 Davey Lab, University Park, PA 16802 ⁸ Department of Physics and Astronomy, College of Charleston,

SC 29424 ⁹ Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile ¹⁰ Millennium Institute of Astrophysics (MAS),

Nuncio Monseñor Sótero Sanz 100, Providencia, Santiago, Chile ¹¹ Space Science Institute, 4750 Walnut Street, Suite 205,

Boulder, Colorado 80301 ¹² Department of Physics, National Institute of Technology,

Nara College, Yamatokohriyama, Nara 639-1080, Japan ¹³ Department of Astronomy, School of Science, Graduate University for Advanced Studies, Mitaka, Tokyo 181-8588, Japan

black holes, 2) they also provide energy and momentum feedback to the interstellar medium of host galaxies and to the intergalactic medium (IGM), and inhibit star formation (e.g., Springel, Di Matteo, & Hernquist 2005), and 3) they may promote metal enrichment of the intergalactic medium (IGM) (e.g., Hamann et al. 1997b; Gabel, Arav, & Kim 2006). These outflowing gases, which are difficult to observe directly, have been detected as absorption lines in the spectra of about 50% of all quasars (e.g., Vestergaard 2003; Wise et al. 2004; Misawa et al. 2007a; Nestor et al. 2008; Muzahid et al. 2013). However, a limitation of these past studies is that they observe the outflowing gas only along a single sightline (i.e., one dimension) toward the nucleus of each quasar, although the absorber's physical conditions probably depend on the location/orientation at which we observe it (e.g., Ganguly et al. 2001; Elvis 2000). Thus, the internal structure of outflowing winds is still largely unknown.

Multiple quasar images, produced by gravitational lensing, provide a unique way to study the outflowing gas along more than one sightline. Lensed quasars with large image separation angles have a higher chance of revealing structural differences in the outflowing winds, especially in the vicinity of the continuum source. In this sense, the quasar images that are lensed by a cluster of galaxies (rather than a single massive galaxy) are very promising targets. Among three such lensed quasars, SDSS J1029+2623 at $z_{em} \sim 2.197$ (Inada et al. 2006; Oguri et al. 2008) is the best target because i) it has the largest lensed quasar image separation ($\theta \sim 22^{\prime\prime}5$) ever observed (see Figure 1 of Inada et al. 2006), and ii) it exhibits absorption features in the blue wings of the C IV, N V, and Ly α emission lines with ejection velocity of $v_{\rm ei} \leq$

1,000 km s⁻¹, which could be the result of outflowing gas moving toward us from the central region. We call them Proximity Absorption Lines (PALs) throughout this paper. We define PALs as a subcategory of Narrow Absorption Lines (NALs) with ejection velocity of $v_{\rm ej} \leq$ 1,000 km s⁻¹. We use this terminology throughout the paper to separate PALs from NALs with larger ejection velocities.

Misawa et al. (2013) obtained high-resolution spectra of the brighter two of the lensed images (images A and B) with the Subaru telescope in 2010 February and found several clear signs that the origin of the PALs is indeed in the outflowing gas. First, they show the signature of partial coverage, which means the absorbers do not cover the background flux source completely. There also exists a clear difference in the absorption profiles between the spectra of images A and B, which can be explained by either of the following two scenarios: (a) time variability of the absorption features over a time scale corresponding to the time delay between the two images (time variation scenario; Chartas et al. $(2007)^{14}$, or (b) a difference in the absorption between the different sightlines of the outflowing wind (*multi-sightline scenario*; Chelouche 2003; Green 2006). However, with a single-epoch observation we cannot distinguish between these scenarios. Misawa et al. (2014b) performed a second observation about four years (1514 days in the observed frame) after the first observation (which is longer than the time delay between images A and B, $\Delta t_{AB} \sim 744$ days), and found that the PALs were nearly stable and that most of the differences between images A and B still remained. This evidence suggests a *multi-sightline scenario* where the absorber's size should be smaller than the physical distance between the sightlines of the lensed images, thus not covering both sightlines. A possible explanation is that there are a number of small *clumpy* clouds in the outflowing stream. Indeed, some of the outflowing gas is expected to consist of small gas clouds $(d_{\text{cloud}} \leq 10^{-3} \text{ pc})$ with very large gas densities $(n_e \geq 10^6 \text{ cm}^{-3}; \text{Hamann et al. 2013; Joshi$ et al. 2014). Furthermore, recent radiation-MHD simulations by Takeuchi et al. (2013) reproduce variable clumpy structures with typical sizes of 20 times the gravitational radius $(R_{\rm g})$, corresponding to $d_{\rm cloud} \sim 5 \times 10^{-4}$ pc, assuming a black-hole mass for SDSS J1029+2623 of $M_{\rm BH} \sim 10^{8.72} {\rm ~M}_{\odot}$ (Misawa et al. 2013).

Such clumpy clouds can be examined more easily through narrow absorption lines (NALs, hereafter) with large offset velocities from the quasar because the corresponding absorbers are (i) probably smaller than the PAL absorbers and (ii) not so crowded in velocity space as the PAL absorbers. Indeed, there are many NALs detected in the spectra of both images A and B of SDSS J1029+2623. Their origin is not only the outflowing wind (*intrinsic* NALs, hereafter) but also cosmologically intervening gas such as foreground galaxies and the IGM (*intervening* NALs, hereafter). Although it has been traditionally believed that many NALs that fall within 5,000 km s⁻¹ of the quasar emission redshift (termed *associated* absorption lines or AALs) are physically associated with the quasar (e.g., Weymann et al. 1979), we can separate intrinsic NALs from intervening ones more effectively by performing partial coverage analysis. In order to form a global picture of the outflowing wind, we need to understand the physical conditions of NAL absorbers (i.e., highly accelerated gas) as well as PAL absorbers (i.e., weakly accelerated gas).

In this paper, we present the results from our new spectroscopic observation of SDSS J1029+2623 taken with the Very Large Telescope (VLT), which enables us for the first time to identify *intrinsic* NALs in a high quality spectrum of this object. In §2, we describe the observations and data reduction. The methods used for absorption line detection and covering factor analysis are outlined in §3. The results and discussion are presented in §4 and §5. Finally, we summarize our results in §6. We use a cosmology with $H_0=70$ km s⁻¹ Mpc⁻¹, $\Omega_m=0.3$, and $\Omega_{\Lambda}=0.7$ throughout the paper.

2. OBSERVATIONS

We acquired high resolution spectra of the brightest two of the three lensed images of SDSS J1029+2623, A and B with V = 18.72 and 18.67 mags, with the VLT using the Ultraviolet and Visual Echelle Spectrograph (UVES) in queue mode (ESO program 092.B-0512(A)). The observations were performed from 2014 January 28 to February 26 (epoch E2, hereafter), which is ~ 4 years after the first observation on 2010 February 10 (epoch E1, hereafter; Misawa et al. 2013), and ~ 2 months before the third observation on 2014 April 4 (epoch E3, hereafter; Misawa et al. 2014b) with Subaru using the High Dispersion Spectrograph (HDS). We used a slit width of 1"2, corresponding to $R \sim 33,000$, while Misawa et al. (2013, 2014b) took $R \sim 30,000$ and 36,000 spectra using Subaru/HDS. The wavelength coverage is 3300–6600 Å in the 390/564 nm setting, which covers the O vI, N v, Si IV, and C IV doublets as well as the $Ly\alpha$ absorption line at $z_{\rm abs} \sim z_{\rm em}$. We also adopted 2×2 pixel binning in both the spatial and dispersion directions to increase the S/N ratio. The total integration time is 26,670s and the final S/N ratio is about 23 pix^{-1} around 4700Å for both images.

We reduced the data to extract the one-dimensional spectra in a standard manner using the UVES Common Pipeline Library (CPL release 6.6). We could not separate the third image (image C, V = 20.63) from image B completely because the typical seeing of our observation (~ 1."0–1."8) was comparable to the separation angle between images B and C ($\theta \sim 1."85$)¹⁵.

Table 1 gives a log of the current observation with VLT/UVES as well as our past observations with Subaru/HDS, in which we list the target name, date of observation, telescope/instrument used, spectral resolution, total exposure time, and signal-to-noise ratio (S/N). The S/N is evaluated around 4700 Å, close to the C IV mini-BAL. In Figure 1, we show normalized spectra over the full wavelength range of our observations for images A and B. These spectra were binned every 0.5Å for display purposes, and the 1 σ errors are also shown.

3. DATA ANALYSIS

¹⁴ The time delay between images A and B is $\Delta t_{AB} \sim 774$ days in the sense of A leading B, while the time delay between images B and C Δt_{BC} is only a few days (Fohlmeister et al. 2013).

¹⁵ On the other hand, a flux contamination from the image C is almost negligible in the image B spectrum because the former is much fainter than the latter.

First, using the line detection code SEARCH, written by Chris Churchill, we detect all absorption features whose confidence level is greater than 5σ in the normalized spectrum of each lensed image. We then identify N v, C IV, and Si IV doublets in the regions from $-1,000 \text{ km s}^{-1}$ to $5,000 \text{ km s}^{-1}$ (N v), from $-1,000 \text{ km s}^{-1}$ to $70,000 \text{ km s}^{-1}$ (C IV), and from $-1,000 \text{ km s}^{-1}$ to $40,000 \text{ km s}^{-1}$ (Si IV) around the corresponding emission lines, with the maximum velocity set in order to avoid the $Ly\alpha$ forest¹⁶. We also search for the Mg II doublet in the whole range of the spectra. Absorption troughs that are separated by nonabsorbed regions are considered to be separate lines. In total, 4 Mg II, 19 C IV, 2 N V, and 7 Si IV doublets are identified in the image A spectrum, while 3 Mg II, 22 C IV, 2 N V, and 7 Si IV doublets are identified in the image B spectrum. The equivalent widths of both blue and red members of doublets are measured for each line by integrating across the absorption profile and these are listed in Table 2. We also searched for 10 single metal lines (O I λ 1302, Si II λ 1190, Si II λ 1193, Si II λ 1260, Si II λ 1527, Al II λ 1671, C II $\lambda 1036$, C II $\lambda 1335$, Si III $\lambda 1207$, and C III $\lambda 1548$) as well as $Lv\alpha$ and $Lv\beta$ and detected about 200 lines at the same redshift as the doublet lines. These are summarized in Table ??. Other single lines or unidentified lines are not shown in Figure 1 and Tables 1-?? even if they are detected at a confidence greater than 5σ .

3.1. dN/dz Analysis

One of the important properties of intrinsic NALs is a number density excess of high-ionization doublets per unit redshift (i.e., dN/dz; Hamann et al. 1997a, and references therein). In order to compare the dN/dz from our spectra with those from our previous study based on 37 quasar spectra (Misawa et al. 2007a), we construct a complete sample including only NALs whose blue doublet members would be detected even in the lowest S/N spectrum in Misawa et al. (2007a). The corresponding lower limits of rest-frame equivalent widths (EWs) are $\mathrm{EW}_{\min}^{\mathrm{rest}} = 0.056 \text{ Å}$, 0.038 Å, and 0.054 Å for C IV, N V, and Si IV, respectively. Here, the values of $\mathrm{EW}_{\min}^{\mathrm{rest}}$ depend on the S/N ratio of the observed spectrum as

$$EW_{\min}^{\text{rest}} = \frac{-U^2 + U\sqrt{U^2 + 4(S/N)^2(M_L^2 M_c^{-1} + M_L)}}{2(S/N)^2(1 + z_{abs})} \times \Delta\lambda(A_L^2 M_c^{-1} + M_L)$$

where U is the confidence level of the EW defined as $\text{EW}/\sigma(\text{EW})$, and M_L and M_C are the numbers of pixels over which the equivalent width and the continuum level are determined (Young et al. 1979; Tytler et al. 1987). Using equation (1), we confirm that the S/N of our VLT spectra is always larger than the required values except for the region between $\lambda_{\text{obs}} \sim 4500 - 4525$ Å (Figure 2). After removing weak NALs with EW^{rest} < EW^{rest}_{min}, we have 12 C IV, 2 N V, and 3 Si IV doublets in image A spectrum and 11 C IV, 2 N V, and 3 Si IV doublets in the image B spectrum. We will call this the "homogeneous" NAL sample, hereafter. Following Misawa et al. (2007a), we also combined multiple NALs lying within 200 km s⁻¹ of each other into a single NAL "system" because clustered lines are probably not independent even if they have a cosmologically intervening origin (e.g., Sargent, Steidel, & Boksenberg 1988).

All identified doublets (including both the homogeneous and inhomogeneous NAL samples) are listed in Table 2, in which multiple NALs within 200 km s⁻¹ of each other are separated by horizontal lines. The table gives the ion name (ion), flux-weighted absorption redshift (z_{abs}) , ejection velocity (v_{ej}) supposing they originate in the outflow, rest frame EWs of blue and red members of the doublet $(EW_{b}^{rest}, EW_{r}^{rest})$, identification number in Figure 1 (ID), reliability class of intrinsic lines (described later), ionization class (described later), and velocity difference from the first doublet at the lowest redshift in each absorption system (Δv). Table ?? summarizes other information including the flux-weighted line width of each system on a velocity scale, $\sigma(v)$, as defined in Misawa et al. (2007a), and other transitions that are detected at the same redshift, as well as the column density $(\log N)$, Doppler parameter, b, and covering factor, $C_{\rm f}$, for each absorption component in the system.

3.2. Partial Coverage Analysis

Among several criteria, i) time variability, ii) partial coverage, and iii) line locking are the most reliable properties to distinguish intrinsic NALs from intervening NALs (Barlow & Sargent 1997; Hamann et al. 1997a, and references therein). When compared with broad intrinsic absorption lines, intrinsic NALs are less likely to vary (Misawa et al. 2014a; Chen et al. 2015) and when they do vary, their variation amplitude is small (Wise et al. 2004; Misawa et al. 2014a). Therefore, the variability criterion does not offer an efficient way of identifying intrinsic NALs.

Partial coverage analysis is quite useful for our spectra because the resolving power is high enough to deblend NALs into multiple components. Using the Voigt profile fitting code MINFIT (Churchill 1997; Churchill et al. 2003), we deblended NALs into 86 and 91 components in images A and B, respectively. We do not include the Si IV NAL at $z_{\rm abs} = 1.8909$ because the blue and red members of the doublet are both blended with other lines. With MINFIT, we fit each NAL profile using the redshift (z), (λ) , column density (log N in cm⁻²), Doppler parameter (b), in km s⁻¹), and covering factor $(C_{\rm f})$ as free parameters. The covering factor $(C_{\rm f})$ is the fraction of photons from the background source that pass through the absorber. If the background source is uniformly bright, then $C_{\rm f}$ also represents the fraction of the background source (i.e., the continuum source and/or broad emission line regions, BELRs) that is occulted by foreground absorbers along our sightline. If $C_{\rm f}$ is less than unity, it is likely that the absorbers are part of a quasar outflow because cosmologically intervening absorbers like substructures in foreground galaxies and the IGM are less likely to have internal structures as small as a size of the background flux sources (e.g., Wampler et al. 1995; Barlow & Sargent 1997). The covering factor is evaluated in an unbiased manner as 1)2

$$C_f = \frac{(R_r - 1)^2}{1 + R_b - 2R_r} , \qquad (2)$$

where R_b and R_r are the residual (i.e., unabsorbed) fluxes

 $^{^{16}}$ Here, we define the velocity offset as positive for blueshifted NALs from the quasar emission redshift that is determined from Mg II emission line (Inada et al. 2006).

of the blue and red members of a doublet in the normalized spectrum (c.f., Hamann et al. 1997b; Barlow & Sargent 1997; Crenshaw et al. 1999). If MINFIT gives unphysical covering factors for some components, e.g., negative or greater than 1, we rerun the code assuming $C_{\rm f} = 1$ only for those components because the $C_{\rm f}$ values are very sensitive to continuum level errors, especially for full coverage doublets (Misawa et al. 2005). In addition to the *fitting* method above, we evaluate $C_{\rm f}$ values for each *pixel* as Ganguly et al. (1999) did (*pixel-by-pixel* method). The fitting results by both methods are shown in Figure 3 and the fit parameters by the fitting method are summarized in Table ??. We have confirmed that the $C_{\rm f}$ values provided by the two methods are in good agreement with each other.

4. RESULTS

4.1. Narrow Absorption Lines (NALs)

Using our NAL sample toward the lensed images of SDSS J1029+2623, we perform several statistical analyses. In these analyses, in order to avoid any possible biases we do not include the two C IV NALs at $z_{\rm abs} \sim 1.9322$ and 1.9788 that are covered in the Subaru spectrum (Misawa et al. 2013) but not covered by our VLT data.

4.1.1. dN/dz analysis

In Table 4, we summarize the number density of homogeneous NAL systems per unit redshift (dN/dz) or per unit velocity offset from the quasar $(dN/d\beta)$ for C iv, N IV, and Si IV, along with the Poisson noise in these quantities (Gehrels 1986). All systems are also classified into one of two categories according to the offset velocity from the quasar; associated absorption lines (AALs) with $v_{\rm ej} \leq 5,000$ km s⁻¹, and non-AALs with $v_{\rm ej} > 5,000$ km s⁻¹, following Misawa et al. (2007a). We found that the dN/dz values for C IV, N V, and Si IV toward this one quasar, SDSS J1029+2623, are larger than the average values in the larger sample of Misawa et al. (2007a) by factors of ~ 3 , ~ 6 , and ~ 2 , respectively, although this enhancement is not statistically significant because of the small number of NALs in our spectra. Because Misawa et al. (2007a) discovered that at least $\sim 20\%$ of C iv NALs originate from winds based on partial coverage analysis, statistically we expect to detect two or more intrinsic C IV NALs among ~ 10 homogeneous C IV NALs in the spectra of images A and B.

4.1.2. Intrinsic or Intervening NALs

High-velocity NALs blueshifted with $v_{\rm ej} > 1000 \,\mathrm{km \, s^{-1}}$ are expected to be outside of the range of velocities where BELs are found, and thus, they absorb mostly continuum light. The existence of partial coverage in high-velocity NALs suggests that the size of absorbers is comparable to or smaller than the continuum source (i.e., $d_{\rm cloud} \leq R_{\rm cont}$). Following Misawa et al. (2007a), we separate all NALs (including those in the inhomogeneous sample) into three classes (classes-A, B, and C) based on partial coverage analysis, where class-A includes NALs most reliably classified as intrinsic while class-C includes NALs that are consistent with full coverage or that cannot be classified. Class-B contains NALs that show line-locking, which is a signature of a radiatively driven outflowing wind and is only detectable if our sightline is approximately parallel to the gas motion, as often seen in NALs (e.g., Benn et al. 2005; Bowler et al. 2014). Class-B also contains systems that have tentative evidence for partial coverage. As a result, 4 C IV NALs (including PALs) are classified as intrinsic (two class-A and two class-B) among the 12 C IV NALs in the homogeneous sample from the spectrum of image A, while 3 C IV NALS are classified as intrinsic (two class-A and one class-B) among 11 NALs in the spectrum of image B. The fraction of intrinsic NALs (27 - 33%) is somewhat larger than the average value of $\sim 20\%$ (Misawa et al. 2007a), although our sample size is small. If we include the inhomogeneous sample, 5 C IV NALs are classified as 3 class-A and 2 class-B among 19 NALs toward image A, while 9 out of 22 C IV NALs are classified into 4 class-A and 5 class-B NALs toward image B. In addition to intrinsic C IV NALS, we detected two class-B Si IV NALS only toward image A in homogeneous sample, and three class-B Si IV NALs in each of the image A and B spectra after including systems from the inhomogeneous sample. It is also noteworthy that we detect a large number of line-locked NALs: 2 C IV and 3 Si IV NALs in image A and 5 C IV and 3 Si IV NALs in image B, while only five systems are line-locked among 138 homogeneous NALs toward 37 quasars (Misawa et al. 2007a). This result strongly suggests that our sightline toward the central source is almost parallel to the outflowing streamline.

4.1.3. Ionization Conditions

The outflowing winds in the vicinity of the flux source are probably more highly ionized than most intervening absorbers due to strong UV radiation from the continuum source, although it depends on the gas density of the absorbers. Indeed, a high ionization state has been used as one indicator of the intrinsic properties of absorbers (Hamann et al. 1997a, and references therein). Broad absorption lines (BALs) are often classified into three categories according to their ionization level: highionization BALs (HiBALs), low-ionization BALs (LoB-ALs), and extremely low-ionization BALs showing Fe II lines (FeLoBALs) (Weymann et al. 1991). A similar classification has also been performed for NALs (Bergeron et al. 1994; Misawa et al. 2007a). Motivated by the literature, we classify our NALs into three categories, based on the detection of absorption lines in low (ionization potential; IP < 25 eV), intermediate (IP = 35 - 50 eV), and high (IP > 60 eV) ionization levels¹⁷. Interestingly, some class-A NAL systems include low ionization lines (see Table 2), which suggests that intrinsic NAL absorbers have multiple phases with different ionization states, as noted in Misawa et al. (2007a).

4.1.4. Similarities in NALs between Sightlines

Here, we present a new method of identifying intrinsic NALs by exploiting our multi-sightline observation. If NALs with similar line profiles are detected at the same redshift toward two sightlines, the corresponding

 $^{^{17}}$ An important caveat here is that we do not necessarily detect all absorption lines because of the detection limits due to line strength, observed wavelength coverage, and line blending (e.g., in Ly α forest).

absorber must have a size larger than the physical distance between the two sightlines. The absorber also cannot have any internal structures on the scale of the sightline separation. Although foreground galaxies and IGM structures also can cover both sight lines (whose physical separation is ~kpc or ~Mpc scale), they usually have significant internal velocity structures on those scales, as often seen in the spectra of lensed quasars (e.g., Ellison et al. 2004). Only intrinsic absorbers can satisfy the requirement of having very similar profiles because their sizes can be larger than the sightline separation (e.g., sub-parsec scale) if they are at a small distance from the flux source (e.g., r < 1 kpc). Based on the ejection velocity distribution of class-A, B, and C NALs (Figure 4), in Figure 5 we summarize the distribution of velocity differences between NALs and PALs in the two sightlines (Δv) for systems within $\Delta v \leq 200 \text{ km s}^{-1}$ between the two sightlines. The distribution is almost uniform up to $\Delta v \sim 200 \text{ km s}^{-1}$ with a clear peak near $\Delta v \sim 0 \text{ km s}^{-1}$. Among eight NAL pairs with $\Delta v \leq 10 \text{ km s}^{-1}$, four are C IV and N V PALs at $z_{\rm abs} \sim z_{\rm em}$, which we will discuss later. The other four NAL pairs are C IV NALs at $z_{\rm abs}$ \sim 1.8909, 2.1349, and 2.1819, and a Si IV pair at $z_{\rm abs}$ ~ 1.8909 . Although the velocity shift is very small for these NAL pairs, their line profiles are clearly different as compared in Figure 6. This means our two sightlines go through different absorbers or different regions of a single absorber. In either event, we cannot conclude that these are intrinsic NALs based only on this analysis. On the other hand, it is intriguing that no NAL pairs that are classified into class-A/B have a common ejection velocity or line profile. For example, the C IV NALs at $z_{\rm abs} = 1.7652$ toward image A and at $z_{\rm abs} = 1.7650$ toward image B are both classified as class-A NALs. Although their ejection velocities are very close each other, their line centers and profiles are obviously different, as shown in Figure 7. We also note that the difference is remarkable in all three epochs (see Figure 8). This means the multi-sightline scenario (i.e., that two sightlines pass through different regions of the outflow) is also applicable for intrinsic NALs with large ejection velocity as well as for PALs as already confirmed in Misawa et al. (2013, 2014b).

4.1.5. Time Variability of NALs

Because of the lower data quality (S/N ~ 10 pixel⁻¹) and narrower effective wavelength coverage of the Subaru/HDS spectra taken in epochs E1 and E3, we cannot monitor NALs for time variability analysis with only a few exceptions. As an example, we compare spectra obtained at three epochs around the strong C IV NALs at $z_{abs} = 1.8909$ –1.9138 (ID = 35–44) in image A and C IV NALs at $z_{abs} = 1.8909$ –1.9119 (ID = 41–48) in image B as shown in Figure 9. These NALs are obviously not variable, which is consistent with the past result that NALs with large ejection velocities are rarely variable (Chen et al. 2015).

4.2. Proximity Absorption Lines (PALs)

The absorption line group at $z_{\rm abs} \sim z_{\rm em}$ with an ejection velocity of $v_{\rm ej} < 1000 \text{ km s}^{-1}$ has already been observed twice in 2010 February (epoch E1) and 2014 April (epoch E3) with Subaru/HDS. Its origin is probably in

the outflowing gas because: i) there is evidence for partial coverage (e.g., there exists a clear residual flux at the bottom of the Ly α and N v absorption lines even though they appear to be saturated.), ii) the profiles are variable, and iii) the profiles show signatures of line-locking (Misawa et al. 2013, 2014b). The most important result from these past observations is that the absorption profiles of the Ly α , N v, and C iv PALs in the spectra of images A and B are clearly different, especially for lines at $v_{\rm ej} < 0 \,\rm km \, s^{-1}$ (see Figure 10). There are at least three possible origins for the difference: a) a micro-lensing effect, b) time delay between the images, and c) different absorber structure along the different sightlines. Among these, the first idea is immediately rejected because the lensed images show a common ratios between the radio, optical, and X-ray fluxes (Ota et al. 2012; Oguri et al. 2012), which is not expected for micro-lensing. We can also reject the time delay effect because the variability is almost negligible between epochs E1 and E3 (Misawa et al. 2014b). A difference in column density between the two sightlines is the only acceptable explanation. To our knowledge, this is the first time that an outflowing wind has been observed along multiple sightlines. Hereafter, we call the PALs at $v_{\rm ej}<0~{\rm km~s^{-1}}$ (showing sightline difference) narrow PALs and those at $v_{\rm ej} > 0 \ {\rm km \ s^{-1}}$ (showing similar line profiles between sightlines) broad PALs (see Figure 10).

Our observation with VLT/UVES not only gives an additional epoch for monitoring the PALs but enables us to study the detailed velocity structure of PALs using higher quality spectra with a S/N ratio double that of the past observations with Subaru. We first confirm the line-locking pattern in velocity plots of the C IV PAL as shown in Figure 11, which was already noted in Misawa et al. (2013). Like the other NAL absorbers, the PAL absorbers also appear to be outflowing almost parallel to our sightline. We also reconfirm that the C $\hat{\rm rv}$ PALs became shallower (i.e., decreasing in equivalent width) over a large velocity range of the profile between epochs E1 and E2/E3 (see Figure 11) at $\Delta t_{\rm rest} \sim 1.3$ years in the quasar rest-frame. However, these lines are almost stable on the shorter time scales of $\Delta t_{\rm rest} \sim 0.05$ years between epochs E2 and E3. The broad spectral coverage of the VLT/UVES spectra allows us to detect for the first time the O VI doublet corresponding to the PAL absorber (see Figure 10). There is also a hint of the Si IV doublet detected, but it appears blended with other lines at lower redshift.

4.2.1. Comparison of Covering Factors

In addition to line profiles and strengths, we also compare covering factors for the clean part of the C IV and N v PALs in the three epochs. The fitting parameters to the PALs in epoch E2 with MINFIT are summarized in Table ??. The total column densities of C IV and N v PALs after summing all Voigt components are $\log(N_{\rm NV}/\rm cm^{-2})$ = 15.62 and $\log(N_{\rm CIV}/\rm cm^{-2})$ = 15.90 in image A and 16.04 and 15.89 in image B spectrum. These values are all consistent with the corresponding values that we measured in epoch E1 (Misawa et al. 2013) with differences of a factor of \leq 2. Because all PALs have small offset velocities and are located on the BELs in the spectra, we should consider the BELRs as well as the continuum source as the background flux source. Because the C IV PAL is partially self-blended (i.e., blending of the blue members of one doublet with the red member of another doublet), we can compare covering factors effectively only for the narrow PALs at $v_{\rm ej} \sim -100-0$ km s⁻¹ (corresponding to $\Delta v \sim 400-500$ km s⁻¹ in Figure 12), for which self-blending is negligible¹⁸.

We show the fitting results for the PALs in epoch E2 in Figure 12 and Table ??. The average covering factors of C IV and N v at $v_{\rm ej} \sim -100-0$ km s⁻¹ (i.e., $\Delta v \sim 400-500$ km s⁻¹ in Figure 12) are $C_{\rm f} = 0.47\pm0.03$ and 0.58 ± 0.05 toward image A, and $C_{\rm f} = 0.23\pm0.02$ and 0.39 ± 0.08 toward image B. These values are all consistent with the corresponding values in epoch E1 toward image A ($C_{\rm f} = 0.47\pm0.05$ and 0.61 ± 0.07) and image B ($C_{\rm f} \sim 0.2$ and 0.35 ± 0.05)¹⁹ and in epoch E3 toward image A ($C_{\rm f} = 0.45\pm0.09$ and 0.54 ± 0.09) and image B ($C_{\rm f} = 0.20\pm0.05$ and 0.34 ± 0.19). Thus, covering factors are not variable at least on a timescale of $\Delta t_{\rm rest} \sim 1.3$ years.

The N v PALs all have larger $C_{\rm f}$ values than that for C IV, which is consistent with past results that higher ionization transitions tend to have larger coverage fractions (e.g., Petitjean & Srianand 1999; Srianand & Petitjean 2000; Misawa et al. 2007a; Muzahid et al. 2015). This result suggests that i) the size (i.e., distance from the central black hole) of the N v BELR is smaller than that of the C IV BELR, and/or ii) the size of the N v absorbers is larger than that of the C IV absorbers. It is also interesting that the PALs in the image A spectrum always have larger $C_{\rm f}$ values than those in the image B spectrum, which is additional evidence for variations in the structure along multiple sightlines.

5. DISCUSSION

5.1. Velocity anisotropies in NAL absorbers

Line-locking is seen in both NALs and PALs, thus we are likely to be observing the quasar almost along the direction of the outflow. Nonetheless, we discovered a velocity difference between the two sightlines (see Table 2). Given the redshifts of the lens ($z_1 = 0.58$; Oguri et al. 2008) and the source ($z_s = 2.197$) the separation angle of the light rays that form images A and B, as seen from the source²⁰, is $\theta' \sim 14$."6. This velocity gradient suggests that the outflowing wind has internal velocity anisotropies on a scale of $\sim 1 \text{ km s}^{-1} \text{ arcsec}^{-1}$ on average. Indeed, the hydrodynamic simulations of Proga & Kallman (2004) show this type of internal velocity variations. A recent radiation-MHD simulation by Takeuchi et al. (2013) also reproduced such velocity variations over a typical spatial scale of ~ 5×10^{-4} pc. This is consistent with the size of NAL absorbers, which is also comparable to the size of the continuum source (~ 2.5×10^{-4} pc; Misawa et al. 2013). However, we should note that the velocity variations in the simulations are found in the inner part of the wind at distances of ~ $50R_{\rm g}$ from the center.

The velocity distribution of intrinsic NALs that are classified into Class-A or B appears to cluster around values of $v_{\rm ej} \sim 59,000, 43,000$, and $29,000 \,{\rm km \, s^{-1}}$, except for two intrinsic NALs at $v_{\rm ej} \sim 49,500 \,{\rm km \, s^{-1}}$ (class-A) and $\sim 8,500 \,{\rm km \, s^{-1}}$ (class-B) in the image B spectrum (Figure 4). These clustering patterns are reminiscent of the filamentary structures obtained by numerical simulations (e.g., Proga, Stone, & Kallman 2000). If these patterns are indeed due to filamentary structures, there should exist some velocity anisotropies within them. Velocity dispersions in the three intrinsic NAL clusters are $\delta v \sim 900, 260, \text{ and } 1200 \,{\rm km \, s^{-1}}$ respectively, which correspond to about 1.6%, 0.6%, and 4.0% of their average ejection velocity.

5.2. Ionization condition in PAL absorbers

The Ly α , C IV, and N V PALs have been monitored at three epochs (E1, E2, and E3) between 2010 February and 2014 April. Between epochs E1 and E2/E3, C IV PALs show a clear variation in their strength (i.e., depth). There are several possible reasons for this time variability: (a) gas motion across our line of sight (e.g., Hamann et al. 2008; Gibson et al. 2008; Muzahid et al. 2015), (b) changes in the ionization state of the absorber (e.g., Hamann et al. 2011; Misawa et al. 2007b), and (c) redirection of photons around the absorber by scattering material (e.g., Lamy & Hutsemékers 2004; Misawa et al. 2010). Among these, the first scenario can be rejected because all absorption components in the C IV PAL vary in concert, which requires the implausible situation in which all clouds cross our sightline simultaneously (c.f. Misawa et al. 2007b). The third scenario is also less likely because it requires a variation in the covering factor while the $C_{\rm f}$ values remain almost stable both in C IV and N V PALs between epochs E1 and E2/E3. Thus, only the scenario involving a change in ionization state deserves further investigation. This scenario is further separated into two variants: i.e., C^{3+} ions are ionized to C^{4+} or they recombine to C^{2+} . Both variants of this scenario can explain the decreasing EW of the C IV PALs. Without knowing an absorber's ionization parameter²¹, we cannot tell which one is more likely. If the latter variant applies, we can place constraints on the electron density and the distance from the ionizing photon source by the same prescription as used in Narayanan et al. (2004), taking the variability time scale as an upper limit to the recombination time. Based on the observation that the C IV PALs vary between epochs E1 and E3 over $\Delta t_{\rm obs} = 1514$ days (i.e., $\Delta t_{\rm rest} = 474 \, {\rm days})^{22}$ we can place a lower limit on the

 $^{^{18}}$ Negative ejection velocity for these components could be due to our underestimation of these values because the emission redshift is determined from broad UV emission lines, as done for this quasar by Inada et al. (2006), which are systematically blueshifted from the systemic redshift that is measured by narrow, forbidden lines (see, e.g., Corbin 1990; Tytler & Fan 1992; Brotherton et al. 1994; Marziani et al. 1996) by about 260 km s⁻¹.

¹⁹ Because no absorption component was used for C IV PAL at $v_{\rm ej} \leq 0 \text{ km s}^{-1}$ in the epoch E1 spectrum (Misawa et al. 2013), we adopt an average line depth for this region as a covering factor.

²⁰ The separation angle as seen by the source is given by $\theta' = (D_{\rm ol}/D_{\rm sl}) \times \theta$, where θ is the observed separation angle of the images (22'.5) and $D_{\rm ol}$ and $D_{\rm sl} = ((1 + z_s)/(1 + z_l)) \times D_{\rm ls}$ are angular diameter distances from the observer to the lens and from the source to the lens, respectively.

²¹ The ionization parameter U is defined as the ratio of hydrogen ionizing photon density (n_{γ}) to the electron density (n_e) .

 $^{^{22}}$ We compare spectra in epochs E1 and E3 instead of epochs E1 and E2 because the E2 observation spans the time range 2014 January 28 to February 26, which gives an additional uncertainty for measuring the variation time scale.

electron density of the absorber as $n_e > 8.7 \times 10^3$ cm⁻³, and an upper limit on the distance from the flux source as r < 620 pc, assuming $U \sim 0.02$, the value at which the C IV and N V ions are close to the optimal ionization states for those elements (e.g., Hamann et al. 1995).

5.3. Global Picture of the Outflow from SDSS J1029+2623

In Table 5, we summarize the physical properties of broad PALs, narrow PALs, and intrinsic NALs. We also present a possible geometry of the outflow along our two sightlines in Figure 13 based on our previous constraints.

First, we see almost the same absorption profiles of $Ly\alpha$, N v, and C v PALs in the two lensed images except for a clear difference in the narrow PALs at $v_{\rm ei} \sim$ -100-0 km s⁻¹. We also confirm that none of the intrinsic NALs have common absorption profiles between the images. These results suggest that the size of the broad PAL absorbers are larger than the projected separation between sightlines at the distance of the absorbers, $r\theta$, while the narrow PAL absorbers and NAL absorbers have sizes smaller than $r\theta$. Such common absorption profiles are, however, observable regardless of the absorber's size, if their distance from the flux source is smaller than the boundary radius $r_{\rm b}$ at which the two sightlines of lensed images become fully separated with no overlap (Misawa et al. 2013). The boundary distance is ~ 3.5 pc for SDSS J1029+2623 if only the continuum source is counted. If we also consider the BELR (whose size is estimated to be $\sim 0.09 \text{ pc})^{23}$ as the background source, $r_{\rm b}$ would be ~1200 pc (see Figure 13).

We can also place constraints on the absorber size based on partial covering analysis. Intrinsic NALs with large ejection velocities have partial coverage, although they absorb only the continuum photons. This means their physical scale is comparable to or smaller than the size of the continuum source, $d_{cloud} \leq 2.5 \times 10^{-4}$ pc. On the other hand, broad PAL absorbers cover almost entirely both the BELR and the continuum source, which means the size of the absorbers as a whole is comparable to or larger than the BELR size, ≥ 0.09 pc. Narrow PAL absorbers may consist of a number of small clumpy clouds because they show partial coverage.

In our VLT/UVES spectra, we detected highionization transitions like O VI and N V in the PAL systems, while the intrinsic NAL systems are in various ionization states, with or without high-ionization transitions, as already noted in the literature (e.g., Misawa et al. 2007a; Ganguly et al. 2013). Some C IV NAL absorbers also show N V transitions, while others do not. Because Ly α and N V in the PAL system are less variable than C IV and because N V has larger covering factor than C IV, the cross-section of Ly α and N V should be larger than that of C IV. PAL absorbers with strong O VI lines are probably located much closer to the ionizing flux source than NAL absorbers, although this conclusion depends on the density of the absorber.

It is well known that both broad and narrow absorption lines at $z_{\rm abs} \sim z_{\rm em}$ tend to vary (Wise et al. 2004; Misawa et al. 2014a), while NALs with large ejection velocities are rarely variable (e.g., Chen et al. 2015). Indeed, the PALs in our spectra are variable, while most of the C IV NALs are probably stable between the three epochs, as we observed for a few strong C IV NALs. Misawa et al. (2014a) suggest that broader absorption lines like the broad PALs can vary mainly due to a change in their ionization state while narrow absorption lines like narrow PALs and NALs vary primarily due to the gas motion transverse to our sightlines. The gas motion scenario could explain why narrow PALs are variable but highvelocity NALs are not. Based on the dynamical model of Murray et al. (1995) and more recent investigations (e.g., Misawa et al. 2005; Hall et al. 2011), the absorbers at larger distance from the center have small transverse (i.e., orbital) velocities compared to their radial velocities. If intrinsic NALs lie at larger distances than narrow PALs, their small transverse velocities would rarely lead to time variability, while the large transverse velocities of narrow PALs can lead to time variability more frequently.

6. SUMMARY

In this study, we performed a spectroscopic observation for images A and B of the gravitationally lensed quasar SDSS J1029+2623, and monitored the absorption profiles in these spectra as well as in the previous two observations. Using high quality spectra taken with VLT/UVES, we detected intrinsic narrow absorption lines (NALs) as well as broader, proximity absorption lines (PALs), and fit models to the line profiles. Based on the results of our multi-sightline spectroscopy, we discuss a possible geometry and internal structure of the outflowing wind along our sightlines. Our main results are as follows.

- We detected 66 NALs, of which 24 are classified as *intrinsic* NALs (physically associated with the quasar) based on partial coverage analysis.
- Class-A and B NALs cluster at $v_{\rm ej} \sim 59,000$, 43,000, and 29,000 km s⁻¹, which is reminiscent of the filamentary structures that are often obtained in numerical simulations.
- Our multiple sightline observation suggests that the size of the broad PAL absorbers are larger than the projected distance between sightlines $(r\theta)$ while the narrow PAL absorbers and intrinsic NAL absorbers have sizes smaller than $r\theta$ if their radial distances are greater than the boundary distance.
- While PAL systems show only high ionization transitions, including O vI, intrinsic NAL systems show a wide range of ionization conditions with and without low ionization transitions like O I, Al II, and Si II, as noted in the literature.
- No class-A/B NALs (i.e., candidates for intrinsic NALs) have common absorption profiles in the two lensed images, which means that the outflow has an internal velocity structure whose typical spatial scale is smaller than the physical distance between the sightlines (i.e., $\leq r\theta$).
- Short-time variation in the PALs is probably due to a change in the ionization state of the gas. If this is the case, we can place a lower limit on the gas

 $^{^{23}}$ This is calculated by Misawa et al. (2013) based on the empirical equation in McLure & Dunlop (2004).

density as $n_e \geq 8.7 \times 10^3 \text{ cm}^{-3}$ and an upper limit on the absorber's distance from the flux source as $r \leq 620$ pc.

• Based on our best knowledge, we present a possible geometry of the outflow along our two sightlines, in which we identify different structures in the outflowing wind that can produce, respectively, broad PALs, narrow PALs, and NALs with large ejection velocities.

For further investigations of the outflow's internal structure, especially in the transverse direction, we should perform the same observations for image C of SDSS J1029+2623. We also aim to observe several lensed quasars with smaller separation angles ($\theta \sim 2''$) by a single massive galaxy to examine ~ 10 times finer internal structure in an outflow, as already mentioned in Misawa et al. (2014b).

We thank the anonymous referee for comments that helped us improve the paper. We would like to thank

- REFF
 Arav, N., Borguet, B., Chamberlain, C., Edmonds, D., & Danforth, C. 2013, MNRAS, 436, 3286
 Benn, C. R., Carballo, R., Holt, J., et al. 2005, MNRAS, 360, 1455
 Barlow, T. A., & Sargent, W. L. W. 1997, AJ, 113, 136
 Bergeron, J., et al. 1994, ApJ, 436, 33
 Blandford, R.D. & Parne, D.G., 1982, MNRAS, 199, 883
 Bowler, R. A. A., Hewett, P. C., Allen, J. T., & Ferland, G. J. 2014, MNRAS, 445, 359
 Brotherton, M. S., Wills, B. J., Steidel, C. C., & Sargent, W. L. W. 1994, ApJ, 423, 131
 Chartas, G., Eracleous, M., Dai, X., Agol, E., & Gallagher, S. 2007, ApJ, 661, 678
 Chen, Z.-F., Gu, Q.-S., Chen, Y.-M., & Cao, Y. 2015, MNRAS, 450, 3904
 Chelouche, D. 2003, ApJ, 596, L43

- 450, 3904
 Chelouche, D. 2003, ApJ, 596, L43
 Churchill, C. W., Vogt, S. S., & Charlton, J. C. 2003, AJ, 125, 98
 Churchill, C. W. 1997, Ph.D. Thesis, Univ. California, Santa Cruz
 Corbin, M. R. 1990, ApJ, 357, 346
 Crenshaw, D. M., Kraemer, S. B., Boggess, A., Maran, S. P., Mushotzky, R. F., & Wu, C.-C. 1999, ApJ, 516, 750
 Gehrels, N. 1986, ApJ, 303, 336
 Dorodnitsyn, A. V. 2009, MNRAS, 393, 1433
 Dunn, J. P., Bautista, M., Arav, N., et al. 2010, ApJ, 709, 611
 Ellison, S. L., Ibata, R., Pettini, M., et al. 2004, A&A, 414, 79
 Elvis, M. 2000, ApJ, 545, 63
 Emmering, R.T., Bladford, R.D., & Shlosman, I., 1992, ApJ, 385, 460 460
- Everett, J. E., 2005, ApJ, 631, 689
- Fohlmeister, J., Kochanek, C. S., Falco, E. E., et al. 2013, ApJ,
- 764, 186 Foltz, C., Wilkes, B., Weymann, R., & Turnshek, D. 1983, PASP, 95, 341
- Gabel, J.R., Arav, N., & Kim, T.-S. 2006, ApJ, 646, 742
 Ganguly, R., Lynch, R. S., Charlton, J. C., et al. 2013, MNRAS, 435, 1233 Ganguly, R., Bond, N. A., Charlton, J. C., et al. 2001, ApJ, 549,
- 133Ganguly, R., Eracleous, M., Charlton, J. C., & Churchill, C. W.
- Galaghy, R., Eracleous, M., Charlton, J. C., & Churchill, C. W 1999, AJ, 117, 2594
 Gibson, R. R., Brandt, W. N., Schneider, D. P., & Gallagher, S. C. 2008, ApJ, 675, 985
 Green, P. J. 2006, ApJ, 644, 733
 Hall, P. B., Brandt, W. N., Petitjean, P., et al. 2013, MNRAS, 424, 292

- 434, 222 Hall, P. B., Anosov, K., White, R. L., et al. 2011, MNRAS, 411,
- Hamann, F., Chartas, G., McGraw, S., et al. 2013, MNRAS, 435, 133
- Hamann, F., Kanekar, N., Prochaska, J. X., et al. 2011, MNRAS, 410, 1957
- Hamann, F., Kaplan, K. F., Rodríguez Hidalgo, P., Prochaska, J. X., & Herbert-Fort, S. 2008, MNRAS, 391, L39
 Hamann, F., & Sabra, B. 2004, AGN Physics with the Sloan Digital Sky Survey, 311, 203

Masamune Oguri, Poshak Gandhi, and Chris Culliton for their valuable comments. We also would like to thank Christopher Churchill for providing us with the MINFIT and SEARCH software packages. The research was supported by the Japan Society for the Promotion of Science through Grant-in-Aid for Scientific Research 15K05020, JGC- \tilde{S} Scholarship Foundation, and partially supported by MEXT Grant-in-Aid for Scientific Research on Innovative Areas (No. 15H05894). CS acknowledges support from CONICYT-Chile through Becas Chile 74140006. FEB acknowledges support from CONICYT-Chile (Basal-CATA PFB-06/2007, FONDECYT Regular 1141218, "EMBIGGEN" Anillo ACT1101) and the Ministry of Economy, Development, and Tourism's Millennium Science Initiative through grant IC120009, awarded to The Millennium Institute of Astrophysics, MAS. JCC and ME acknowledge support from the National Science Foundation through award AST-1312686.

REFERENCES

- Hamann, F., Barlow, T. A., Junkkarinen, V., & Burbidge, E. M. 1997a, <u>ApJ</u>, 478, 80 Hamann, F., Barlow, T. A., & Junkkarinen, V. 1997b, ApJ, 478,
- 87 Hamann, F., Barlow, T. A., Beaver, E. A., et al. 1995, ApJ, 443,
- 606
- Inada, N., Oguri, M., Morokuma, T., et al. 2006, ApJ, 653, L97 Jannuzi, B. T., Hartig, G. F., Kirhakos, S., et al. 1996, ApJ, 470, L11

- Joshi, B. T., Hatug, C. P., Hinako, J., et al. 1556, ApJ, 476, L11
 Joshi, R., Chand, H., Srianand, R., & Majumdar, J. 2014, MNRAS, 442, 862
 Konigl, A., & Kartje, J. F. 1994, ApJ, 434, 446
 Kurosawa, R. & Proga, D. 2009, ApJ, 693, 1929
 Lamy, H., & Hutsemékers, D. 2004, A&A, 427, 107
 Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., & Moles, M. 1996, ApJS, 104, 37
 McLure, R. J., & Dunlop, J. S. 2004, MNRAS, 352, 1390
 Misawa, T., Charlton, J. C., & Eracleous, M. 2014a, ApJ, 792, 77
 Misawa, T., Inada, N., Oguri, M., et al. 2014b, ApJ, 794, L20
 Misawa, T., Kawabata, K. S., Eracleous, M., Charlton, J. C., & Kashikawa, N. 2010, ApJ, 719, 1890
 Misawa, T., Charlton, J. C., Eracleous, M., Ganguly, R., Tytler, D., Kirkman, D., Suzuki, N., & Lubin, D. 2007a, ApJS, 171, 1
 Misawa, T., Eracleous, M., Charlton, J. C., & Kashikawa, N.

- D., KHKMAH, D., SUZUKI, N., & LUDIN, D. 2007a, ApJS, 171,
 Misawa, T., Eracleous, M., Charlton, J. C., & Kashikawa, N. 2007b, ApJ, 660, 152
 Misawa, T., Eracleous, M., Charlton, J.C., & Tajitsu, A., 2005, ApJ, 629, 115

- ApJ, 629, 115
 Murray, N., Chiang, J., Grossman, S. A., & Voit, G. M., 1995,
 ApJ, 451, 498
 Muzahid, S., Srianand, R., Charlton, J., & Eracleous, M. 2015,
 arXiv:1509.07850
- arXiv:1509.07850
 Muzahid, S., Srianand, R., Arav, N., Savage, B. D., & Narayanan, A. 2013, MNRAS, 431, 2885
 Narayanan, D., Hamann, F., Barlow, T., Burbidge, E.M., Cohen, R.D., Junkkaribeb, V., & Lyons, R., 2004, ApJ, 601, 715
 Nestor, D., Hamann, F., & Rodriguez Hidalgo, P. 2008, MNRAS, 386, 2055

- 386, 2055
 Nomura, M., Ohsuga, K., Wada, K., Susa, H., & Misawa, T. 2013, PASJ, 65,
 Oguri, M., Schrabback, T., Jullo, E., et al. 2012, arXiv:1209.0458
 Oguri, M., Ofek, E. O., Inada, N., et al. 2008, ApJ, 676, L1
 Ota, N., Oguri, M., Dai, X., et al. 2012, arXiv:1202.1645
 Petitjean, P. & Srianand, R., 1999, A&A, 345, 73
 Proga, D., & Kallman, T. R. 2004, ApJ, 616, 688
 Proga, D., Stone, J. M., & Kallman, T. R., 2000, ApJ, 543, 686
 Sargent, W.L.W., Boksenberg, A., & Steidel, C.C., 1988, ApJS, 68, 539

- 68, 539

- 68, 539
 Springel, V., Di Matteo, T., & Hernquist, L. 2005, ApJ, 620, L79
 Srianand, R. & Petitjean, P., 2000, A&A, 357, 414
 Takeuchi, S., Ohsuga, K., & Mineshige, S. 2013, PASJ, 65, 88
 Tytler, D., & Fan, X.-M. 1992, ApJS, 79, 1
 Tytler, D., Boksenberg, A., Sargent, W. L. W., Young, P., & Kunth, D., 1987, ApJS, 64, 667

- Vestergaard, M., 2003, ApJ, 599, 116
 Wampler, E. J., Chugai, N. N., & Petitjean, P. 1995, ApJ, 443, 586
 Weymann, R. J., Morris, S. L., Foltz, C. B., & Hewett, P. C. 1991, ApJ, 373, 23
 Weymann, R. J., Williams, R. E., Peterson, B. M., & Turnshek, D. A. 1979, ApJ, 234, 33
- Wise, J. H., Eracleous, M., Charlton, J. C., & Ganguly, R. 2004, ApJ, 613, 129
 Young, P. J., Sargent, W. L. W., Boksenberg, A., Carswell, R. F., & Whelan, J. A. J., 1979, ApJ, 229, 891

Target	Obs Date	Instrument	R	$T_{ m exp}$ (sec)	${ m S/N^a} \ ({ m pix}^{-1})$	$\operatorname{Reference}^{b}$
SDSS J1029+2623 A	2010 Feb 10 2014 Jan 28 – Feb 3 2014 Apr 4	Subaru/HDS VLT/UVES Subaru/HDS	$30000 \\ 33000 \\ 36000$	$14400 \\ 26670 \\ 11400$	$13 \\ 23 \\ 14$	$\begin{array}{c}1\\2\\3\end{array}$
SDSS J1029+2623 B	2010 Feb 10 2014 Feb 4 – 26 2014 Apr 4	Subaru/HDS VLT/UVES Subaru/HDS	30000 33000 36000	$14200 \\ 26670 \\ 11400$	$13 \\ 23 \\ 14$	$\begin{array}{c}1\\2\\3\end{array}$

TABLE 1 Log of Observations

 $^{\rm a}$ Signal to noise ratio at $\lambda_{\rm obs} \sim 4700 {\rm \AA}.$ $^{\rm b}$ References. (1) Misawa et al. 2013, (2) This paper, (3) Misawa et al. 2014b.

		Δv^e cm s ⁻¹)		$^{0.0}_{+19.8}$		0.0 + 125.1	+172.0	00	+22.9 +126.2	0.0 + 103.0			+66.8		00	+108.6	0.0	0.0	0.0	0.0	0.0 + 10.4	+51.8 +51.8	$^{0.0}_{+72.5}$	$^{0.0}_{+31.0}$	0.0 0.0	+30.9 +41.2	+236.9 +267.8	¢	+29.0	0.0 + 125.0	+57.5	+143.9 0.0 0.0
		(k									M	ult	i-S	igh	ntli	ine	Ob	ose:	rvat	tion	of	NA	Ls									11
		Class-2 ⁶		Γ	Γ	Г	Ц	Ħ	Ē	Н	H	ΞÞ	4	Η	111 1	ПЛ	Η		HI		LIH		LIH	LIH		LIH			H	LIH	HI	H
		Class-1 ^c		C3	C1	C1	Бq	4 0	D2	C2,H	B2	55	77	A2	60	BI	A2		C2,H	B2	C2,H C2.H		B2	С1,H С1,H B2		C2,H C1,H			Ы,Н	C1.H	C2	С1,н C2
		Ð		31, 32	49,50	67,68	210	19.19	12,10	16,20	18,22	25,26	67,17	29,30	<i>3</i> 6 66	33,30 $34,37$	35,38		41, 43	5,14	$6,17 \\ 42.44$		8,19	$^{9,21}_{45,46}_{11,23}$		$15,24 \\ 47,48$		2	51,52	53,54	55,57	59,60
	ge B	$\mathrm{EW}^{\mathrm{rest}b}_{\mathrm{r}}$ $(\mathrm{\AA})$		0.052 ± 0.004	1.444 ± 0.008	0.092 ± 0.005	<u>000 0+960 0</u>	200-01-020-00	761.0	0.144 ± 0.002	0.014 ± 0.002	0.081 ± 0.003	U.U40±U.UU2	0.032 ± 0.002	0 01 1 0 000	0.066 ± 0.002	0.022 ± 0.002		≤ 0.501	0.119 ± 0.004	$\leq 0.383 \leq 0.592$	1	0.019 ± 0.002	$\begin{array}{c} 0.142 \pm 0.002 \\ 0.363 \pm 0.006 \\ 0.013 \pm 0.002 \end{array}$		≤ 0.197 0.679 ± 0.009			0.110 ± 0.004	0.470 ± 0.003	0.014 ± 0.002	0.032±0.002
	Imag	EW_{b}^{resta} (Å)		0.095 ± 0.004	1.689 ± 0.008	$0.186 {\pm} 0.005$	0.035+0.009	200.0T00700	0.100±0.10	≤ 0.284	0.021 ± 0.002	0.133 ± 0.004	U.U04±U.UU2	0.042 ± 0.002	0 010 1 0 000	0.094 ± 0.002	0.041 ± 0.002		$0.456 {\pm} 0.006$	0.121 ± 0.004	0.380 ± 0.005 < 0.669	1	0.032 ± 0.002	$\begin{array}{c} 0.162 \pm 0.003 \\ 0.457 \pm 0.005 \\ 0.023 \pm 0.002 \end{array}$		≤ 0.248 0.837 ± 0.009			0.209±0.004	0.552 ± 0.003	0.029 ± 0.002	0.183±0.002 0.073±0.002
		${v_{\rm ej} \over {\rm (km \ s^{-1})}}$		190276	171003	141157	RDDDR	00200	09490	58699	58350	55033 ±1090	67010	49509	195.06	43590 43496	43250		30088	30087	29737 29723		29413	$28991 \\ 28964 \\ 28917$		$27949 \\ 27946$		0100	8460	7223	6585 222	65UU 5886
		$z_{\rm abs}$		0.5124	0.6731	0.9184	1 6085	1 6140	1.0149	1.6221	1.6253	1.6556	1.0324	1.7065	41041	1.7627 1.7627	1.7650		1.8909	1.8909	$1.8944 \\ 1.8945$		1.8975	$1.9016 \\ 1.9019 \\ 1.9024 $		$1.9118 \\ 1.9119$		1004	2.1081	2.1209	2.1276	2.1285
		Ion		Mg II	Mg II	Mg II	C IV			C IV	C IV		~ 1 ^	C IV	i H	22 00	C IV		C IV	Si IV	Si IV C IV		Si IV	Si IV C IV Si IV		Si IV C IV			CIV	C IV	C IC	GIV
		Class-2 ^d	L	Г		L			HI	Н			LIH		H		111	HIT				LIH	H		LIH		LIH	LI	HI	LH	Н	Η
-		Class-1 ^c	C1	C1		C3	CI		$^{ m B2,H}_{ m C2}$	B2,H			C2,H		С2,Н		64	C1.H		D2,D		$^{ m C2,H}_{ m B2}$	C1,H C2		С1,Н С1,Н		С3,Н В2,Н	C2	C2	C2	CI	CI
		Ð	27, 29	28,30		61,63	62, 64		$^{8,11}_{9,12}$	16,18			23, 24	04 00	25,26		91.90	35,37	- С - - - - - - - - - - - - - - - - - -	0,10		$36,39 \\ 6,14$	38,40 7,17		$\frac{41,43}{13,19}$		$\frac{42,44}{15,20}$	21, 22	45, 46	47,48	49,50	51,52
	Ā	$\mathrm{EW}^{\mathrm{rest}b}_{\mathrm{r}}$	$0.495 {\pm} 0.008$	0.145 ± 0.004		0.120 ± 0.002	1.366 ± 0.006		$0.297\pm0.003 \le 0.033$	0.190 ± 0.007			0.185 ± 0.007	0000	0.053 ± 0.004		600 07 260 0	0.309 ± 0.003		ZU.200		$\begin{array}{c} 0.318 \pm 0.004 \\ \leq 0.464 \end{array}$	$\begin{array}{c} 0.146 \pm 0.005 \\ 0.013 \pm 0.002 \end{array}$		0.448 ± 0.007 0.318 ± 0.004		0.359 ± 0.009 0.205 ± 0.005	0.019 ± 0.003	0.074 ± 0.002	0.024 ± 0.001	0.114 ± 0.001	0.046 ± 0.002
	Image /	${ m EW}_{ m b}^{ m resta}$	$0.604{\pm}0.008$	0.221 ± 0.005		0.193 ± 0.002	1.472 ± 0.006		$\begin{array}{c} 0.308{\pm}0.003 \\ 0.047{\pm}0.003 \end{array}$	≤ 0.512			0.498 ± 0.005	0 100 0 002	0.493 ± 0.005		0.047±0.003	0.355+0.004		CUU.UTELOU.U		0.413 ± 0.004 0.137 ± 0.003	$\begin{array}{c} 0.193 \pm 0.004 \\ 0.029 \pm 0.003 \end{array}$		0.572 ± 0.008 0.439 ± 0.004		$0.484\pm0.009 \le 0.461$	0.041 ± 0.002	0.130 ± 0.002	≤ 0.065	0.145 ± 0.001	0.093 ± 0.003
		$\mathop{\rm km}\limits^{v_{\rm ej}} \mathop{\rm s}\nolimits^{-1})$	190431	190265		141245	141116		59472 59378	58601			50965	10000	47932		reeer	30091		oonne		$29686 \\ 29685$	29343 29342		$27981 \\ 27987$		27745 27718	24880	8426	7348	6643	5884
		$z_{\rm abs}$	0.5111	0.5125		0.9176	0.9187		$1.6151 \\ 1.6160$	1.6230			1.6930	- 104 D	1.7212		1 7650	1.8909	0000	1.0909		$1.8949 \\ 1.8949$	1.8982 1.8982		1.9115 1.9115		1.9138 1.9141	1.9420	2.1084	2.1196	2.1270	2.1349
		Ion	Mg II	Mg II		Mg II	Mg II		CIV C	C IV			C IV	Ċ	CIV		μ _C			A1 10		C IV Si IV	C IV Si IV		C IV Si IV		C IV Si IV	Si IV	CIV	C IV	C IV	C IV

TABLE 2 Absorption Systems with Doublet Lines

						Absor	PTION SYST	FABLE 2 TEMS WITH	DOUBLEI	r Lines					12	
C IV Si IV	$2.1800 \\ 2.1815$	$\begin{array}{c} 1604 \\ 1458 \end{array}$	$\begin{array}{c} 0.011 \pm 0.001 \\ 0.115 \pm 0.006 \end{array}$	$\begin{array}{c} 0.008 {\pm} 0.001 \\ 0.041 {\pm} 0.005 \end{array}$	53,55 33,34	C1 C1	HI	i								0.0 + 141.5
C IV	2.1819	1416	0.425 ± 0.004	≤ 0.297	54,56	C1,H		CIV	2.1818	1428	0.298 ± 0.002	0.209 ± 0.002	61,62	С1,Н	HIT	$^{+169.7}_{+179.2}$
								Si IV	2.1821	1403	0.088 ± 0.002	0.062 ± 0.003	39,40	Cl		+198.0
								CIV	2.1897	686	0.848 ± 0.003	≤ 1.405	63,64	A1,H	Η	0.0
C IV	2.1898	676	0.902 ± 0.003	≤ 1.253	57,58	A1,H	Η									+9.4
NV	2.1900	657	0.945 ± 0.004	< 0.882	1, 3	A1,H										+28.2
				I				ΝV	2.1900	658	0.888 ± 0.004	≤ 0.717	1, 3	A1,H		+28.2
N V	2.1955	141	≤ 1.494	1.341 ± 0.004	2, 4	A1,H	Н									0.0
								N V	2.1955	141	≤ 1.231	1.201 ± 0.003	2, 4	A1,H	Н	0.0
								C IV	2.1957	122	1.569 ± 0.003	1.224 ± 0.003	65,66	A1,H		+18.8
C IV	2.1958	113	≤ 1.590	1.303 ± 0.003	59,60	A1,H										+28.2
a Rest-fr	me equivale	ant width	of blue compon	ant of doublet 1	line											

^b Rest-frame equivalent width of red component of doublet line.

^c Reliability class of intrinsic lines based on partial coverage analysis, defined in Misawa et al. (2007a). If an equivalent width is large enough, we classify the doublet as homogeneous

sample with a mark of "H". ^d Ionization class of absorption system with high (H; IP > 60 eV), intermediate (l; IP = 35 – 50 eV), and/or low (L; IP < 25 eV) ionization transitions. ^e Velocity difference from the first doublet in each absorption *system*.

					Μ	ulti-Sightl	ine Ob	oservat	ion of NA	.Ls					13
BLE 3 tow Absorption Components	Other Ions ^f	age A	Mg I A2853, Fe II A2600 (Fe II A2383)	Mg I A2853, Fe II A2600 (Fe II A2344, Fe II A2383)	Mg I A2853, Fe II A2600 (Al III A1863)	Mg I A2853, Fe II A2344,2600 (Al III A1863)	Si IV λ1394,1403			Si II A1527, Al II A1671 (O I A1302, Si II A1260, C II A1335, Si IV A1394)	Lyα	Ly α (O I A1302, Si II A1207, Si IV A1394)	Ly α, Si II λ1527, Al II λ1671, (Si II λ1190,1192,1260, C II λ1335, Si III λ1207)	Ly α, Si II	Lyа (С II А1335, Si III А1207, N V А1239,1243)
TAF RS OF NARR	$C_{\mathrm{f}}{}^{e}$	Ima	$\begin{array}{c} 0.93 \substack{+0.07\\ 0.93 \substack{-0.06\\ 1.00 \substack{+0.06\\ -0.06\\ 0.19 \substack{+0.10\\ -0.09\\ 1.00 \end{array}}$	$\frac{1.00}{0.83\substack{+0.46\\-0.28}}$	1.00	$\begin{array}{c} 1.00\\ 1.00\\ 1.00\\ 0.99\substack{+0.03\\-0.03}\end{array}$	$1.00\substack{+0.04\\-0.04}$	$0.42\substack{+0.08\\-0.08}$	1.00 1.00 1.00	1.00 1.00	1.00 1.00	$0.58\substack{+0.06\\-0.06}$	$0.96\substack{+0.05\\-0.05}$	1.00 1.00 1.00	1.00
PARAMETE	$\sigma(v)/b^d \ ({ m km~s^{-1}})$		$\begin{array}{c} 47.3\\ 5.1\pm0.2\\ 10.5\pm0.5\\ 37.1\pm7.2\\ 7.6\pm0.3\end{array}$	$17.9 \\ 6.0\pm0.4 \\ 13.3\pm4.2$	$\frac{14.1}{4.7\pm0.2}$	$\begin{array}{c} 68.1 \\ 4.1\pm0.3 \\ 16.9\pm0.2 \\ 22.0\pm0.5 \end{array}$	$26.0 \\ 8.7 \pm 0.1$	$13.1 \\ 10.3 \pm 1.1$	38.3 13.0 ± 0.8 5.4 ± 0.6 9.7 ± 3.0	58.4 39.0 ± 1.8 17.9 ± 0.7	$^{41.9}_{11.1\pm0.7}$ $^{9.2\pm0.7}_{9.2\pm0.7}$	$10.0 \\ 6.1 \pm 0.5$	30.8 20.9 ± 0.5	$\begin{array}{c} 27.1 \\ 2.8 \pm 0.9 \\ 8.8 \pm 3.1 \\ 11.2 \pm 0.4 \end{array}$	36.5 7.7 ± 3.0
LINE	$\frac{\log N}{(\mathrm{cm}^{-2})}$		$\begin{array}{c} 12.96 {\pm} 0.06 \\ 13.51 {\pm} 0.05 \\ 13.69 {\pm} 0.33 \\ 12.60 {\pm} 0.02 \end{array}$	$\frac{12.69\pm0.02}{12.70\pm0.27}$	12.95 ± 0.02	$\begin{array}{c} 12.50 {\pm} 0.02 \\ 14.03 {\pm} 0.02 \\ 14.94 {\pm} 0.07 \end{array}$	17.33 ± 0.06	13.66 ± 0.15	$\begin{array}{c} 13.84 {\pm} 0.02 \\ 13.93 {\pm} 0.06 \\ 12.88 {\pm} 0.09 \end{array}$	$\frac{13.69\pm0.02}{13.52\pm0.02}$	$\frac{13.25\pm0.02}{13.16\pm0.02}$	13.71 ± 0.11	14.83 ± 0.05	$\begin{array}{c} 12.48 {\pm} 0.11 \\ 12.49 {\pm} 0.14 \\ 13.56 {\pm} 0.02 \end{array}$	12.85 ± 0.26
	$v_{\rm ej}{}^c$ (km s ⁻¹)		$\begin{array}{c} 190431\\ 190464\\ 190426\\ 190423\\ 190423\\ 190408\end{array}$	$\begin{array}{c} 190265\\ 190269\\ 190261\end{array}$	$\frac{141245}{141246}$	$\begin{array}{c} 141116\\ 141190\\ 141146\\ 141094\end{array}$	59472 59473	59378 59379	58601 58617 58590 58538	50965 50991 50925	$\begin{array}{c} 47932 \\ 47955 \\ 47908 \end{array}$	43224 43223	$30091 \\ 30090$	$\begin{array}{c} 30088 \\ 30121 \\ 30111 \\ 30081 \end{array}$	29686 29728
	$z_{\mathrm{abs}}{}^{b}$		$\begin{array}{c} 0.5111\\ 0.5108\\ 0.5112\\ 0.5112\\ 0.5113\\ 0.5113\end{array}$	$\begin{array}{c} 0.5125\\ 0.5125\\ 0.5126\\ 0.5126\end{array}$	$0.9176 \\ 0.9176$	$\begin{array}{c} 0.9187\\ 0.9181\\ 0.9185\\ 0.9189\\ 0.9189\end{array}$	1.6151 1.6151	$1.6160 \\ 1.6160$	$\begin{array}{c} 1.6230\\ 1.6229\\ 1.6231\\ 1.6236\\ 1.6236\end{array}$	$\begin{array}{c} 1.6930 \\ 1.6928 \\ 1.6934 \end{array}$	$\begin{array}{c} 1.7212 \\ 1.7209 \\ 1.7214 \end{array}$	$1.7652 \\ 1.7652$	1.8909 1.8909	$\begin{array}{c} 1.8909\\ 1.8906\\ 1.8907\\ 1.8910\end{array}$	$1.8949 \\ 1.8945$
	$\lambda_{obs}{}^a$ (Å)		4225.6	4229.5	5376.1	5365.3	4048.7	4050.1	4067.7	4169.3	4212.9	4281.1	4475.7	4029.3	4481.8
	Ion		Mg II	Mg II	Mg II	Mg II	C IV	C IV	C IV	C IV	C IV	C IV	C IV	Si IV	C IV

			Multi-Sightline Observation of NALs			15
350RPTION COMPONENTS		(Si 111 Å1207)	Si IV λ1394, O VI λ1032,1038 Si IV λ1394, O VI λ1032,1038		λ2600	$\lambda 2853$, Fe II $\lambda 2344,2383,2600$
TABLE 3 [ARROW A1) 1.23 1.23	1.10 Ly α	Lya, Lya,	Image B	Fe II .29	Mg I 10 109 108
ERS OF N	$1.00 \\ 0.72 \substack{+0 \\ -0}$	$\begin{array}{c} 0.88\substack{+0\\-0}\\1.00\\1.00\end{array}$	$\begin{array}{c} 1.00\\ 0.38\pm 0\\ 0.38\pm 0\\$		$0^{+}69.0$	$\begin{array}{c} 0.31 \substack{+ 0 \\ - 0 \end{array} \\ 0.92 \substack{+ 0 \\ - 0 \end{array}$
e Paramet	19.5 ± 4.4 3.0 ± 1.1	$\begin{array}{c} 34.0\\ 16.5\pm1.7\\ 29.1\pm0.3\\ 6.0\pm0.3\end{array}$	$\begin{array}{c} 400.1\\ 25.6\pm0.8\\ 11.6\pm0.6\\ 8.6\pm1.3\\ 11.1.6\pm0.6\\ 8.6\pm1.3\\ 11.1.1.1\\ 11.1\pm1.1\\ 12.8\pm0.5\\ 12.8\pm0.0\\ 11.2\pm1.0\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm0.2\\ 12.8\pm1.2\\ 12$		11.7 11.3 ± 0.9	78.2 2.4 ± 1.9 7.5 ± 1.7
LINI	$12.54{\pm}0.07\\12.94{\pm}0.25$	$\begin{array}{c} 13.69{\pm}0.09\\ 13.89{\pm}0.01\\ 13.61{\pm}0.01 \end{array}$	13.24±0.01 14.49±0.07 14.27±0.23 13.77±0.20 14.30±0.02 14.49±0.02 14.49±0.03 13.31±0.01 14.63±0.02 14.88±0.03 14.48±0.02 14.48±0.02 14.48±0.02 14.48±0.02 14.48±0.02 13.91±0.03 14.28±0.03 14.55±0.02 13.90±0.01 14.88±0.03 14.55±0.02 13.90±0.01 14.88±0.03 14.55±0.02 13.90±0.01 14.88±0.03 14.55±0.02 13.90±0.01 14.88±0.03 14.55±0.02 13.90±0.01 15.45±0.33 15.21±0.05 13.56±0.01 15.21±0.05 13.56±0.01		12.68 ± 0.18	13.02 ± 1.03 13.49 ± 0.14
	$1427 \\ 1395$	$1416\\1436\\1409\\1394$	$\begin{array}{c} 347\\ 878\\ 878\\ 878\\ 878\\ 831\\ 795\\ 773\\ 693\\ 693\\ 693\\ 611\\ -11\\ -11\\ -124\\ 61\\ -11\\ -124\\ 61\\ -11\\ -12\\ 888\\ 834\\ 889\\ 834\\ 889\\ 834\\ 889\\ 888\\ 834\\ 889\\ 888\\ 889\\ 888\\ 888\\ 888\\ 888\\ 88$		$190276 \\ 190275$	$\frac{171003}{171070}$
	2.1818 2.1822	$\begin{array}{c} 2.1819 \\ 2.1817 \\ 2.1820 \\ 2.1822 \end{array}$	2.1937 2.1881 2.1886 2.1886 2.1889 2.1909 2.1909 2.1914 2.1961 2.1961 2.1981 2.1981 2.1966 2.1981 2.1981 2.1966 2.1981 2.19666 2.19666 2.19666 2.19666 2.196666 2.19666666 2.196666666666666666		$0.5124 \\ 0.5124$	$\begin{array}{c} 0.6731 \\ 0.6725 \\ 0.6726 \end{array}$
		4926.3	4944.5 3956.5		4229.3	4678.6
		CIV	C IV g N V h		Mg II	Mg II

16					М	isawa	et al.							
SLE 3 ow Absorption Components				(Si IV A1394,1403)						Lyα	Lya, Si II λ 1527 (Si II λ 1260, C II λ 	Lyα	Ly α (Si III λ 1207)	$Ly\alpha(Si III \lambda 1207)$
TAB RS OF NARR	$\begin{array}{c} 1.00 \substack{+0.21\\-0.18\\1.00 \substack{+0.06\\-0.06\\0.98 \substack{+0.06\\-0.06\\1.00\\1.00 \substack{+0.06\\-0.05\\-0.14\\0.55 \substack{+0.14\\-0.14\\-0.14\end{array}} \end{array}$	$\begin{array}{c} 0.29 \substack{+\infty \\ -0.32 \\ 0.68 \substack{+0.18 \\ -0.13 \\ 1.00 \end{array}}$	$0.44\substack{+0.10\\-0.10}$	$\frac{1.00}{0.71^{+0.06}_{-0.06}}$	1.00	$0.28\substack{+0.08\\-0.08}$	$0.85\substack{+0.07\\-0.06}$	1.00	$0.46\substack{+0.07\\-0.06}$	$0.15\substack{+0.11\\-0.11}$	$\begin{array}{c} 0.83\substack{+0.07\\-0.06}\\ 0.17\substack{+\infty\\-0.07\end{array}\end{array}$	$0.36\substack{+0.12\\-0.10}$	1.00 1.00 1.00	
e Parametei	$\begin{array}{c} 16.5 \pm 18.7 \\ 8.1 \pm 2.4 \\ 12.0 \pm 0.3 \\ 3.2 \pm 9.0 \\ 16.5 \pm 0.4 \\ 20.7 \pm 5.0 \\ 17.6 \pm 0.3 \\ 6.0 \pm 2.0 \end{array}$	33.1 9.1 ± 2.0 9.6 ± 1.0 11.8 ± 0.7	$11.2 \\ 8.0 \pm 0.9$	$\begin{array}{c} 20.6 \\ 2.5 \pm 4.5 \\ 9.2 \pm 0.4 \end{array}$	$19.0 \\ 13.4 \pm 0.4$	13.7 6.0 \pm 0.9	22.1 14.7±0.5	$14.7 \\ 11.4\pm0.5$	11.7 7.8 ± 0.8	$11.1 \\ 4.5 \pm 25.6$	20.9 7.6 ± 0.7 22.0 ± 16.2	15.8 11.7±1.4	$\begin{array}{c} 48.6\\11.4\pm0.8\\25.0\pm2.3\\10.4\pm6.0\\11.3\pm2.2\end{array}$	38.2
LINI	$\begin{array}{c} 12.83\pm 0.52\\ 13.03\pm 0.24\\ 13.27\pm 0.01\\ 14.29\pm 9.79\\ 13.29\pm 0.01\\ 13.67\pm 0.16\\ 13.867\pm 0.16\\ 13.38\pm 0.44\\ 13.38\pm 0.44\end{array}$	$\begin{array}{c} 12.45{\pm}0.59\\ 12.65{\pm}0.12\\ 12.22{\pm}0.02 \end{array}$	13.57 ± 0.18	12.38 ± 0.08 14.04 ± 0.06	14.12 ± 0.02	13.57 ± 0.24	13.82 ± 0.06	13.42 ± 0.01	13.68 ± 0.14	14.17 ± 9.99	$\frac{13.78\pm0.07}{13.92\pm0.37}$	$13.64 {\pm} 0.20$	$\begin{array}{c} 13.42 \pm 0.03 \\ 14.31 \pm 0.06 \\ 13.15 \pm 0.59 \\ 13.00 \pm 0.08 \end{array}$	
	$\begin{array}{c} 171047\\ 171032\\ 171018\\ 171008\\ 171008\\ 170994\\ 170980\\ 170955\\ 170947\end{array}$	$\begin{array}{c} 141157\\ 141186\\ 141158\\ 141158\\ 141142\end{array}$	60206 60206	59493 59491 59487	58699 58699	58350 58348	55033 55033	51029 51027	49509 49508	43596 43596	43496 43503 43479	43250 43249	30088 30140 30084 30055 30027	30087
	$\begin{array}{c} 0.6727\\ 0.6729\\ 0.6730\\ 0.6731\\ 0.6732\\ 0.6733\\ 0.6733\\ 0.6735\\ 0.6735\\ 0.6736\end{array}$	$\begin{array}{c} 0.9184 \\ 0.9181 \\ 0.9184 \\ 0.9185 \end{array}$	$\frac{1.6085}{1.6085}$	$\begin{array}{c} 1.6149 \\ 1.6150 \\ 1.6150 \end{array}$	$1.6221 \\ 1.6222$	1.6253 1.6253	$\frac{1.6556}{1.6556}$	$1.6924 \\ 1.6925$	1.7065 1.7065	1.7617 1.7617	$\begin{array}{c} 1.7627\\ 1.7626\\ 1.7628\\ 1.7628\end{array}$	1.7650 1.7650	$\begin{array}{c} 1.8909\\ 1.8904\\ 1.8910\\ 1.8913\\ 1.8913\\ 1.8915\end{array}$	1.8909
		5364.4	4038.4	4048.4	4066.3	4064.5	4111.4	4168.4	4190.2	4275.7	4277.1	4280.7	4475.7	4029.3
		Mg II	C IV	C IV	C IV	C IV	C IV	C IV	C IV	C IV	C IV	C IV	GIV	${ m Si}{ m IV}^i$

				Mult	i-Sightlin	ie Obs	ervation	of NAL	s				1′
ROW ABSORPTION COMPONENTS	Lyα, Si II λ1527, Al II λ1671 (Si II λ1190,1193,1260, C II λ1335, Si III λ1207)	Lya, Si II À1527, Al II À1671 (Si II À1190,1193,1260, C II À1335, Si III À1207)	Lya, C IV A1548,1551 (Si 11 A1260, Si 111 A1207)	Lyα, Si II λ 1527, Al II λ 1671 (Si II λ 1190, 1193, 1260, C II λ 1335, Si III λ 1207)	Lyα, Si II λ1527, Al II λ1671 (Si II λ1190,1193,1260, C II λ1335, Si III λ1207)	Lya, C IV A1548,1551	Ly α, Si II λ1527, Al II λ1671 (О I λ1302, Si II λ1190,1193,1260, Si III λ1207)	Ly α, Si II λ1527, Al II λ1671 (О I λ1302, Si II λ1190,1193,1260, Si III λ1207)	Lya, Si IV $\lambda 1394, 1403$ (Si III $\lambda 1207)$	Lyα, О I А1302Si II А1260, С II А1335, Si IV А1394,1403 (Si II А1207)	Lyα	Lya, Si IV A1394,1403	Lyα
rs of Narr	1.00 1.00 1.00	0001 1000 1000 1000 1000 1000 1000 100	$0.50\substack{+0.16\\-0.13}$	$0.92\substack{+0.05\\-0.05}$	$0.96\substack{+0.07\\-0.07\\0.97\substack{+0.08\\-0.07\\-0.07\end{aligned}}$	$0.31\substack{+0.10\\-0.09}$	$\frac{1.00}{0.96\substack{+0.04\\-0.04}}$	$\frac{1.00}{0.98^{+0.09}_{-0.09}}$	$\begin{array}{c} 1.00\\ 0.70^{+0.46}\\ 1.00\\ 0.70^{-0.25}\\ 0.09\\ \end{array}$	$\begin{array}{c} 0.96\substack{+0.04\\-0.04\\0.92\substack{+0.04\\-0.04\\1.00\end{array}\end{array}$	1.00	$0.97\substack{+0.04\\-0.04}$	1.00
PARAMETE	$\begin{array}{c} 76.4\\22.5\pm0.4\\10.1\pm2.1\\11.3\pm0.7\\12.6\pm0.9\end{array}$	$\begin{array}{c} 75.2\\ 16.9\pm1.9\\ 18.7\pm1.4\\ 7.7\pm1.7\\ 16.6\pm1.2\\ 31.2\pm5.3\end{array}$	$\frac{11.2}{7.4\pm0.9}$	18.5 12.8±0.3	52.4 21.6 ± 0.5 12.5 ± 0.4	$ \begin{array}{c} 11.1 \\ 6.8 \pm 1.2 \end{array} $	$\begin{array}{c} 26.7 \\ 13.5\pm0.2 \\ 18.0\pm1.9 \end{array}$	88.7 34.6 ± 1.8 38.3 ± 0.9	$\begin{array}{c} 62.0\\ 28.8\pm1.9\\ 8.3\pm1.1\\ 15.7\pm0.9\\ 9.0\pm1.0\end{array}$	$\begin{array}{c} 55.1 \\ 14.3 \pm 0.4 \\ 10.1 \pm 0.4 \\ 13.2 \pm 0.2 \end{array}$	$22.9 \\ 15.0 \pm 1.5$	$19.4 \\ 12.8 \pm 0.2$	$32.8 \\ 10.1 \pm 0.8$
LINE	$egin{array}{c} 13.87 \pm 0.02 \ 12.33 \pm 0.08 \ 12.92 \pm 0.02 \ 12.92 \pm 0.02 \ 12.70 \pm 0.03 \end{array}$	$\begin{array}{c} 14.14\pm0.09\\ 14.51\pm0.05\\ 13.23\pm0.14\\ 14.25\pm0.03\\ 13.76\pm0.03\\ 13.76\pm0.07\end{array}$	13.07 ± 0.20	13.87 ± 0.04	$\frac{14.51\pm0.03}{13.87\pm0.05}$	13.19 ± 0.25	$\frac{13.38\pm0.01}{13.67\pm0.04}$	$\frac{13.73\pm0.02}{15.09\pm0.04}$	$\begin{array}{c} 13.12\pm0.02\\ 13.27\pm0.21\\ 13.06\pm0.02\\ 13.64\pm0.07\\ 13.64\pm0.07\end{array}$	$\begin{array}{c} 14.42 {\pm} 0.03 \\ 14.04 {\pm} 0.03 \\ 14.35 {\pm} 0.02 \end{array}$	12.93 ± 0.03	14.22 ± 0.02	12.83 ± 0.03
	29737 29771 29700 29675 29629	29723 29753 29753 29768 29682 29638	29413 29414	$28991 \\ 28992$	28964 28988 28918	28917 28918	27949 27964 27933	27946 28034 27921	8460 8508 8486 8432 8416	7223 7261 7225 7180	6585 6588	6500 6500	$5886 \\ 5913$
	$\begin{array}{c} 1.8944 \\ 1.8940 \\ 1.8947 \\ 1.8950 \\ 1.8954 \end{array}$	$\begin{array}{c} 1.8945\\ 1.8939\\ 1.8942\\ 1.8946\\ 1.8946\\ 1.8949\\ 1.8953\end{array}$	1.8975 1.8975	$1.9016 \\ 1.9016$	$\begin{array}{c} 1.9019 \\ 1.9017 \\ 1.9024 \end{array}$	$1.9024 \\ 1.9023$	$\begin{array}{c} 1.9118 \\ 1.9117 \\ 1.9120 \end{array}$	$\begin{array}{c} 1.9119 \\ 1.9110 \\ 1.9121 \end{array}$	$\begin{array}{c} 2.1081\\ 2.1076\\ 2.1078\\ 2.1084\\ 2.1085\end{array}$	$\begin{array}{c} 2.1209\\ 2.1205\\ 2.1209\\ 2.1214\\ \end{array}$	$2.1276 \\ 2.1276$	2.1285 2.1285	$2.1349 \\ 2.1346$
	4034.0	4488.7	4038.4	4044.2	4492.7	4045.2	4058.4	4508.1	4811.9	4831.8	4842.1	4843.5	4853.4
	Si IV	C IV	Si IV	Si IV	C IV	Si IV	Si IV	C IV	CIV	CIV	C IV	C IV	C IV

18				Misawa et al.
LE 3 M Absorption Components		Lya, Si 11 A1260, Si IV A1394,1403 (Si 11 A1193, Si 111 A1207)	Lya, Si 11 A1260, Si IV A1394,1403 (Si 11 A1193, Si 111 A1207)	Lya, Si II A1260, Si IV A1394,1403 (Si II A1193, Si III A1207) Lya, Si II A1260, Si IV A1394,1403 (Si II A1193, Si III A1207)
TAB: RS OF NARRO	1.00	$\begin{array}{c} 1.00\\ 1.00\\ 1.00\\ 1.00\\ 0.97\substack{+0.03\\-0.03}\end{array}$	${\begin{array}{*{20}c} 0.38 + 0.07 \\ 0.38 - 0.07 \\ 0.77 + 0.10 \\ 0.77 - 0.10 \end{array}}$	$\begin{array}{c} 0.22 +0.02\\0.73 \substack{+0.03\\0.62 \substack{+0.03\\0.62 \substack{+0.03\\0.62 \substack{+0.03\\0.61 \substack{+0.0$
e Paramete	$20.4{\pm}1.2$	$\begin{array}{c} 48.5\\ 16.1\pm3.4\\ 13.9\pm0.5\\ 13.3\pm0.6\\ 14.1\pm0.2\end{array}$	$12.1 \\ 5.2 \pm 1.3 \\ 7.2 \pm 0.6$	$\begin{array}{c} 393.5\\ 39.55\pm 3.5\\ 35.5\pm 3.5\\ 45.5\pm 3.0\\ 23.1\pm 1.5\\ 18.4\pm 0.8\\ 29.2\pm 2.6\\ 22.7\pm 0.3\\ 17.4\pm 0.9\\ 11.3\pm 1.0\\ 29.5\pm 2.0\\ 7.6\pm 1.7\\ 17.1\pm 1.4\\ 25.5\pm 1.5\\ 4.02.7\\ 13.4\pm 9.3\\ 9.3\pm 1.7\\ 17.1\pm 1.4\\ 25.5\pm 2.0\\ 17.9\pm 0.4\\ 31.1\pm 0.5\\ 9.5\pm 1.5\\ 9.5\pm 1.5\\ 24.5\pm 1.7\\ 51.6\pm 1.0\\ 29.5\pm 2.2\\ 29.5\pm 29.5\\ 29.5$
LIN	13.12 ± 0.02	$\begin{array}{c} 12.66\pm0.08\\ 13.30\pm0.02\\ 12.91\pm0.02\\ 14.17\pm0.01 \end{array}$	13.69 ± 0.31 13.16 ± 0.11	$\begin{array}{c} 14.45\pm0.07\\ 14.45\pm0.06\\ 14.45\pm0.06\\ 14.27\pm0.06\\ 13.85\pm0.39\\ 14.50\pm0.01\\ 14.50\pm0.05\\ 14.18\pm0.05\\ 14.18\pm0.02\\ 14.19\pm0.04\\ 14.19\pm0.04\\ 14.19\pm0.04\\ 14.19\pm0.04\\ 14.19\pm0.02\\ 14.22\pm0.28\\ 14.19\pm0.02\\ 14.45\pm1.01\\ 14.60\pm0.07\\ 14.85\pm0.02\\ 14.85\pm0.02\\ 14.55\pm0.02\\ 14.55\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 15.26\pm0.02\\ 14.16\pm0.13\\ 14.16\pm0.13$
	5871	$\begin{array}{c} 1428 \\ 1503 \\ 1471 \\ 1431 \\ 1431 \\ 1406 \end{array}$	$1403 \\ 1411 \\ 1399$	$\begin{array}{c} 348\\ 840\\ 840\\ 689\\ 689\\ 614\\ 614\\ 613\\ 204\\ 167\\ 120\\ 203\\ 120\\ 80\\ 80\\ 80\\ 80\\ 857\\ 759\\ 857\\ 759\\ 857\\ 701\\ 613\\ 856\\ 857\\ 701\\ 857\\ 701\\ 856\\ 857\\ 701\\ 857\\ 701\\ 857\\ 701\\ 856\\ 857\\ 701\\ 856\\ 857\\ 701\\ 856\\ 857\\ 701\\ 86\\ 857\\ 701\\ 86\\ 86\\ 86\\ 86\\ 86\\ 86\\ 86\\ 86\\ 86\\ 86$
	2.1350	$\begin{array}{c} 2.1818\\ 2.1810\\ 2.1814\\ 2.1818\\ 2.1818\\ 2.1821\end{array}$	2.1821 2.1820 2.1821	2.1937 2.1937 2.1885 2.1896 2.1913 2.1913 2.1913 2.1949 2.1956 2.1970 2.1966 2.1970 2
		4926.1	4435.0	4944.5 3956.2
		C IV	Si IV	C IV ^j

TABLE 3 Line Parameters of Narrow Absorption Components

- ^a Wavelength of flux-weighted line center.
 - ^b Redshift of flux weighted line center.
- ^c Ejection velocity from quasar emission redshift.
- ^d Flux-weighted line width $(\sigma(v))$ or Doppler parameter (b).
 - ^e Covering factor.
- f Other single lines that are detected in the system. Lines in parenthesis are in Ly α forest.
- g Two C IV PALs at $z_{abs} \sim 2.1898$ and 2.1958 are fitted together as a single absorption profile whose flux-weighted redshift is $z_{abs} \sim 2.1937$. ^h Two N V PALs at $z_{abs} \sim 2.1900$ and 2.1955 are fitted together as a single absorption profile whose flux-weighted redshift is $z_{abs} \sim 2.1937$.
 - ^j Two C IV PALs at $z_{abs} \sim 2.1897$ and 2.1957 are fitted together as a single absorption profile whose flux-weighted redshift is $z_{abs} \sim 2.1937$. ⁱ We cannot fit this Si IV NAL because both members of the doublet are blending with other lines.
- ^k Two N V PALs at $z_{abs} \sim 2.1900$ and 2.1955 are fitted together as a single absorption profile whose flux-weighted redshift is $z_{abs} \sim 2.1935$.

Misawa et al.

TABLE 4	
STATISTICAL PROPERTIES OF POISSON SYSTEMS OF NALS	

			C IV			N V			Si IV	
	-	AAL^a	non- AAL^b	total	AAL^{a}	non-AAL ^b	total	AAL^a	non- AAL^b	total
Path length	$\delta z^c \ \delta eta^c$	$\begin{array}{c} 0.05 \\ 0.02 \end{array}$	$0.56 \\ 0.19$	$\begin{array}{c} 0.61 \\ 0.21 \end{array}$	$\begin{array}{c} 0.05 \\ 0.02 \end{array}$	$\begin{array}{c} 0.01 \\ 0.00 \end{array}$	$\begin{array}{c} 0.07 \\ 0.02 \end{array}$	$\begin{array}{c} 0.05 \\ 0.02 \end{array}$	$\begin{array}{c} 0.37\\ 0.12\end{array}$	$\begin{array}{c} 0.42\\ 0.14\end{array}$
Image A	N^d dN/dz dN/deta	$\substack{\substack{3^{+2.9}_{-1.6}\\57^{+55}_{-31}\\177^{+172}_{-96}}$	$9^{+4.1}_{-2.9}\\16^{+7.3}_{-5.3}\\47^{+22}_{-16}$	$12^{+4.6}_{-3.4}\\20^{+7.4}_{-5.6}\\58^{+22}_{-17}$	$2^{+2.6}_{-1.3} \\ 38^{+50}_{-24} \\ 118^{+155}_{-76}$	$0^{+1.8}_{-0}\\0^{+132}_{-0}\\0^{+409}_{-0}$	$\begin{array}{r}2^{+2.6}_{-1.3}\\30^{+39}_{-19}\\93^{+123}_{-60}\end{array}$	$\begin{array}{c}0^{+1.8}_{-0}\\0^{+35}_{-0}\\0^{+108}_{-0}\end{array}$	$3^{+2.9}_{-1.6}_{8.1^{+7.9}_{-4.4}}_{25^{+24}_{-14}}$	$\begin{array}{r}3^{+2.9}_{-1.6}\\7.1^{+6.9}_{-3.9}\\22^{+21}_{-12}\end{array}$
Image B	N^d dN/dz dN/deta	$\begin{array}{r} 3^{+2.9}_{-1.6} \\ 57^{+55}_{-31} \\ 177^{+172}_{-96} \end{array}$	$8^{+4.0}_{-2.8} \\ 14^{+7.1}_{-4.9} \\ 42^{+21}_{-15}$	$11^{+4.4}_{-3.3}\\18^{+7.2}_{-5.3}\\53^{+21}_{-16}$	$2^{+2.6}_{-1.3} \\ 38^{+50}_{-24} \\ 118^{+155}_{-76}$	$0^{+1.8}_{-0} \\ 0^{+132}_{-0} \\ 0^{+409}_{-0}$	$2^{+2.6}_{-1.3}\\30^{+39}_{-19}\\93^{+123}_{-60}$	$0^{+1.8}_{-0} \\ 0^{+35}_{-0} \\ 0^{+108}_{-0}$	$3^{+2.9}_{-1.6} \\ 8.1^{+7.9}_{-4.4} \\ 25^{+24}_{-14}$	$\begin{array}{r} 3^{+2.9}_{-1.6} \\ 7.1^{+6.9}_{-3.9} \\ 22^{+21}_{-12} \end{array}$
$M07a^e$	$\frac{dN/dz}{dN/d\beta}$	14. 54.	$6.6 \\ 24.$	$6.9 \\ 25.$	$5.2 \\ 21.$	$\begin{array}{c} 0.0\\ 0.0\end{array}$	4.6 18.	3.2 13.	3.7 14.	$3.7 \\ 14.$

 $\overline{\rm a}$ Associated absorption lines with $v_{\rm ej} \leq 5000 \ {\rm km \ s^{-1}}$ from the quasar emission redshift.

^b Non-associated absorption lines with $v_{\rm ej} > 5000 \text{ km s}^{-1}$ from the quasar emission redshift. ^c Total redshift and speed intervals considered in the determination of dN/dz and $dN/d\beta$.

^d Number of Poisson systems (i.e., groups of NALs that lie within 200 km s⁻¹ of each other). ^e dN/dz and $dN/d\beta$ values from Misawa et al. (2007a).

		TAB	LE 5		
Properties	OF	PALS	AND	INTRINSIC	NALS

	$\begin{array}{c} \mathrm{PAL}^{a}\\ \mathrm{(broad)} \end{array}$	PAL^b (narrow)	NAL	$reference^{c}$
Absorber's Transverse Size ^d Ionization Condition ^e Radial Velocity ^f (v_{ej}) Rotational Velocity ^g (v_{rot}) Variability Origin of Variability Covering Factor (C_f) Background flux source	$ \begin{array}{c} \geq r\theta \\ \text{high} \\ \text{small} \\ \text{large} \\ \text{frequently} \\ \text{ionization condition} \\ \sim 1 \\ \text{BELR} \end{array} $	$\leq r\theta$ high small large frequently gas motion < 1 BELR	$ \begin{array}{c} \leq \mathrm{r}\theta \\ \mathrm{low}-\mathrm{high} \\ \mathrm{large} \\ \mathrm{small} \\ \mathrm{very} \ \mathrm{rare}^h \\ \mathrm{gas} \ \mathrm{motion} \ (\mathrm{if} \ \mathrm{any}) \\ < 1 \\ \mathrm{continuum} \ \mathrm{source} \end{array} $	$1 \\ 1, 2, 3 \\ 1 \\ 7, 8, 9 \\ 4, 5, 6 \\ 5 \\ 1 \\ 1$

^a PALs at $v_{\rm ej} > 0$ km s⁻¹ with common absorption profiles in both sightlines.

^b PALs at $v_{ej} > 0$ km s⁻¹ with common absorption promes in both signtimes. ^c References. (1) This paper, (2) Misawa et al. (2007a), (3) Ganguly et al. (2013) (4) Wise et al. (2004), (5) Misawa et al. (2014a), (6) Chen et al. (2015), (7) Murray et al. (1995), (8) Misawa et al. (2005), (9) Hall et al. (2011)

^d This depends on whether sightline difference is observed (i.e., $\leq r\theta$) or not (i.e., $\geq r\theta$). ^e Low, intermediate, and high ionization transitions are defined in Section 4.1.3.

^f Ejection velocity assuming $z_{\rm em} = 2.197$. ^g Absorbers with small orbital radius tend to have larger rotational velocity (see Section 5.3).

^h Variation probability is very low for high ejection velocity NALs (Chen et al. 2015), although we confirmed this trend only for a few strong C IV NALs.



FIG. 1.— Normalized spectra and their 1σ errors for images A and B of SDSS J1029+2623, taken with VLT/UVES, after sampling every 0.5Å for display purpose only.



FIG. 1.— Continued.



FIG. 1.— Continued.



FIG. 2.— Signal-to-Noise (S/N) ratio of the image A and B spectra (blue histograms) and the minimum S/N ratio that is necessary for detecting homogeneous sample lines for N V, C IV, and Si IV, respectively (red lines from top to bottom; see Section 3.1).



FIG. 3.— Results of partial coverage analysis applied to all C IV, N V, Si IV, and Mg II NALs detected in images A and B of SDSS J1029+2623, except for the PALs whose results are presented in Figure 12. The horizontal axis denotes the relative velocity from the flux-weighted center of the system (Δv) while the vertical axis is the normalized flux. The first two panels show the profiles of the blue and red members of a doublet (blue and red histograms) with the model profile produced by MINFIT superposed (dashed line). The positions of the narrow components are marked with upward arrows in the bottom of each panel. The bottom panel shows the covering factors with their 1 σ error bars, measured for each narrow component by MINFIT (green circles) or for each pixel (black dots). If we had to assume $C_{\rm f} = 1$ for some components because of unphysical $C_{\rm f}$ values derived from MINFIT (see Section 3.2), we do not plot them.

















FIG. 4.— Ejection velocity distribution of class-A (red line), B (blue line), and C (black line) NALs and PALs along the sightlines to the images A and B. All Mg II NALs are located outside of the range of the velocity plot. Each of the red lines at $v_{\rm ej} \sim 0$ km s⁻¹ surrounded by a solid rectangle includes both C IV and N V PALs, respectively. Three clustering regions are surrounded by dotted rectangles.



FIG. 5.— Distribution of velocity offset (Δv) between PALs (red histogram) and class-A/B NALs (blue histogram) in the two different images (image A and B) that match each other within $\Delta v \leq 200$ km s⁻¹, as discussed in Section 4.1.4.



FIG. 6.— Comparison of spectra and their 1 σ errors around C IV NALs at $z_{\rm abs} \sim 1.8909$ (upper left), 2.1349 (upper right), 2.1819 (lower left) and Si IV NAL at $z_{\rm abs} \sim 1.8909$ (lower right) in image A (blue histogram) and B (red histogram) spectra for systems that have a small relative velocity between the images.



FIG. 7.— Same as Figure 6 but for class-A C IV NALs at $z_{abs} \sim 1.7652$ in image A (blue histogram) and at $z_{abs} \sim 1.7650$ in image B (red histogram). The latter corresponds to the system center ($\Delta v = 0 \text{ km s}^{-1}$) in this plot.



FIG. 8.— Comparison of C IV NALs at $z_{abs} = 1.7652$ in image A (top panel) and at $z_{abs} = 1.7650$ in image B (bottom panel). Black, red, and blue histograms denote spectra in epoch E1, E2, and E3, respectively. The positions of blue and red member of the doublet are marked with down arrows and vertical dashed lines. A strong feature at $\lambda \sim 4284$ Å in the bottom panel is C IV 1551 at $z_{abs} = 1.7627$. Regions at $\lambda \leq 4281$ Å are not covered by the Subaru/HDS spectra in epochs E1 and E3 because they are located at the edge of the CCD.



FIG. 9.— Comparison of spectra and their averaged 1σ errors around C IV NALs at $z_{abs} \sim 1.8909-1.8982$ in image A (upper left), 1.9115–1.9138 in image A (upper right), 1.8909–1.9019 in image B (lower left) and 1.9119 in image B (lower right) taken in epoch E1 (black histogram), E2 (red histogram), and E3 (blue histogram).



FIG. 10.— Comparison of normalized spectra around Ly α , Si IV, C IV, N V, and O VI PALs detected in the images A (blue) and B (red). The horizontal axis is the offset velocity from the quasar emission redshift, and the vertical axis is the normalized flux. The histograms above the zero flux line are 1σ flux errors. Due to heavy blending with other lines at lower redshift the existence of the Si IV doublet may not be present. The O VI 1032 locate at the edge of the observed spectrum. Absorption features at $v_{ej} < 0 \text{ km s}^{-1}$ in the shaded areas are narrow PALs, while the other features at $v_{ej} > 0 \text{ km s}^{-1}$ are broad PALs.



FIG. 11.— Comparison of C IV, N V, and Ly α PALs at $z_{abs} \sim z_{em}$ in images A and B, taken in epochs E1 (black), E2 (red), and E3 (blue). These spectra are scaled to each other by applying a least-square method for unabsorbed wavelength regions at both sides of the PALs. The vertical axis denotes an arbitrary flux because we did not perform absolute flux calibration without flux standard stars. The gray histograms above the zero flux line are 1σ flux error after taking averages of those in three observing epochs. Double-headed arrows in the bottom of the first two panels mark the velocity ranges of the blue and red members of the C IV and N V doublets.

CIV1548 NV1239 E3 A CIV E3 ----0.5 0.5 0 0 CIV155 NV¹24 ບ້ ి 0.5 Flux, 0.5 Flux, c 0 0.5 0.5 0 -500 500 -500 0 0 500 $\Delta v \ (km \ s^{-1})$ $\Delta v (km s^{-1})$ CIV154 NV123 0.5 0.5 0 0 CIV155 NV1245 చ్ ບ້ Flux, 0.5 Flux, 0.5 0 0 0.5 0.5 0 0 <u>t</u>tt -500 500 -500 500 $\Delta v \ (km \ s^{-1})$ $\Delta v \ (km \ s^{-1})$

FIG. 12.— Same figures as Figure 3, but for C IV and N V PALs at $z_{abs} \sim z_{em}$. We adopt z = 2.1927 as a system center only for the purpose of displaying these velocity plots. Because C IV PALs are self-blended, we fit the profiles of the blue and red members of the doublet simultaneously by multiplying the contributions from them (Misawa et al. 2007b). Spectral regions in the top two panels, for which the profiles of the blue and red members of a doublet (blue and red histograms) and the model profile (dashed line) do not match, suffer from self-blending.



FIG. 13.— Possible geometry of the outflowing wind along our two sightlines toward the lensed quasar SDSS J1029+2623, based on our observations in three epochs. The gray filled circle corresponds to the broad PAL absorber with the minimum acceptable size, while blue and red filled circles are C IV and N V absorbing regions in clumpy clouds as candidates for narrow PAL absorber and intrinsic NALs. The boundary radius of the continuum source (3.5 pc) and BELR (1200 pc) are marked with dashed lines. The NAL absorber's distance is not necessarily larger than 1200 pc, although it is at least larger than the distance of the PAL absorbers. We adopt 14.''6 as the separation angle between two sightlines seen from the source, while Misawa et al. (2013, 2014b) used 22.''5 assuming it is very similar to that seen from us. See Section 5.3 for details.